

Disclaimer: Information contained in the report addresses environmental conditions only and is not the official South Florida Water Management District operations recommendation or decision.

M E M O R A N D U M

TO: John Mitnik, Assistant Executive Director, Executive Office Staff

FROM: SFWMD Staff Environmental Advisory Team

DATE: December 4, 2024

SUBJECT: Weekly Environmental Conditions for Systems Operations

Summary

Weather Conditions and Forecast

Dry and stable air mass associated with a high pressure system will spread over the SFWMD Wednesday morning, inhibiting any rainfall while temperatures remain below normal. After the high moves into southwestern Atlantic Ocean, low-level winds will allow for a gradual warm-up to commence after a chilly morning, the coldest so far this fall season. The dry weather from early in the week will continue through the Thursday. Next, a very strong upper-air disturbance will bring a cold front to north Florida by midnight on Friday and push into the Kissimmee Valley Friday morning, to Lake Okeechobee during the afternoon, the southern third of the SFWMD by the evening, and clear the area entirely by midnight on Saturday. The orientation of the upper-air disturbance will ensure that no rainfall occurs with it. This will occur, even though there will be a surge of greater moisture funneled from the Gulf of Mexico across Florida ahead of it. After the frontal passage, the air mass's strong stability and lack of moisture will help to extend the dry streak through the weekend while temperatures are near or slightly below seasonal averages. By Sunday afternoon, a warming trend will begin, with some increase of shallow moisture by Monday. However, strong atmospheric stability will preclude any rainfall from occurring. For the week ending next Tuesday morning, there is high confidence that no total SFWMD rainfall will occur, nor is any forecast across the SFWMD, which should it verify, will make it 18 days since the last measurable rainfall. Moreover, by that time, only a tenth of an inch of total rainfall will have occurred across the SFWMD over the last month. For the week-2 period (*10-16 December*), all signs point to a very warm weather pattern with below-normal rainfall likely continuing.

Kissimmee

Minor releases were made in the last week from East Lake Toho and Lake Toho to keep the lakes at their regulation schedules. Weekly average discharge on December 1, 2024, was 1,500 cfs and 1,400 cfs at S-65 and S-65A, respectively. Mean weekly water depth on the Kissimmee River floodplain decreased by 0.04 feet to 0.74 feet. The weekly average concentration of dissolved oxygen in the Kissimmee River increased from 4.6

mg/L the previous week to 6.2 mg/L, which is above both the potentially lethal level of 1.0 mg/L and the stressful level of 2.0 mg/L (**Figure KB-6**).

Lake Okeechobee

Lake Okeechobee stage was 14.45 feet NAVD88 (15.76 ft NGVD29) on December 1, 2024, which was 0.15 feet lower than the previous week and 0.34 feet lower than a month ago. Average daily inflows (excluding rainfall) decreased from 1,700 cfs the previous week, to 1,420 cfs. Average daily outflows (excluding evapotranspiration) increased from 2,550 cfs the previous week to 3,670 cfs. The most recent non-obscured satellite image from November 28, 2024, suggests minimal bloom activity on Lake Okeechobee.

Estuaries

Total inflow to the St. Lucie Estuary averaged 289 cfs over the past week with most of the flow coming from the Tidal Basin. Mean salinities increased at all three sites within the estuary. Salinity in the middle estuary was within the optimal range (5-10) for adult eastern oysters.

Total inflow to the Caloosahatchee Estuary averaged 2,083 cfs over the past week. Salinities continued to decrease at all sites within the estuary. Fort Myers surface sensor is currently offline and not reporting this week. Salinities were in the optimal range (0-10) for tape grass in the upper estuary. Salinities were in the optimal range (10-25) for adult eastern oysters at Cape Coral and Shell Point, and in the upper stressed range at Sanibel.

Stormwater Treatment Areas

For the week ending Sunday, December 1, 2024, 3,300 ac-ft of Lake Okeechobee water was delivered to the FEBs/STAs. The total amount of Lake releases sent to the FEBs/STAs in WY2025 (since May 1, 2024) is approximately 81,100 ac-feet. The total amount of inflows to the STAs in WY2025 is approximately 969,000 ac-feet. STA cells are near target stage. STA-1E Central Flow-way is offline for construction activities. Operational restrictions are in effect in STA-1E Western Flow-way, STA-1W Northern Flow-way, STA-2 Flow-ways 2 and 4, and STA-3/4 Eastern Flow-way for vegetation management activities. This week, if LOSOM recommends Lake releases to the WCAs and conditions allow, releases will be sent to STA-1E, STA-2, or STA-3/4.

Everglades

Rates of stage change over the week were categorized as fair or good last week with exception to northern WCA3A which is experiencing excessive recession rates. Water depths throughout the Everglades Protection Area (EPA) have decreased, more substantially in the northern portions of the WCAs and throughout Big Cypress and southern Everglades National Park (ENP). Conditions in WCA-3A North continue to be drier relative to the rest of that basin which may have negative implications for the upcoming wading bird nesting season. Recession rates of up to 0.12 feet per week are considered protective of good foraging conditions. Large numbers of birds are already foraging along the coast allowing for early nesting and coastal colony formation and the relatively wet conditions across the landscape could produce critical foraging habitat in over-drained higher elevation areas during the nesting period if recession rates are appropriate. Average stage in Taylor Slough decreased last week and remains above the recent average (since 2017) for this time of year. Average stage in Taylor Slough decreased last week and remains above the recent average (since 2017) for this time of year. Average salinity increased throughout Florida Bay last week with greatest increases occurring in the western offshore region. Salinities in the eastern region of Florida Bay are now in the 25th percentile of the Inter-Quartile Range (IQR) while the central and western region now fall just below the IQR. Florida Bay MFL metrics remain well outside thresholds of harm.

Biscayne Bay

Total inflow to Biscayne Bay averaged 299 cfs, and the previous 30-day mean inflow averaged 338 cfs. The seven-day mean salinity was 28.9 at BBCW8 and 23.6 at BBCW10, both within the ideal salinity range for estuarine organisms in this region (salinity less than 35). Data were provided by Biscayne National Park.

Supporting Information

Kissimmee Basin

Upper Kissimmee

On December 1, 2024, mean daily lake stages were 57.0 feet NAVD88 (at schedule) in East Lake Toho, 54.0 feet NAVD88 (55.0 feet NGVD29 and at schedule) in Lake Toho (although a discrepancy caused by the datum conversion causes stage to appear higher in the hydrograph in **Figure KB-2**), and 50.4 feet NAVD88 (1.4 feet below the Increment 1 temporary deviation schedule) in Lakes Kissimmee-Cypress-Hatchineha (KCH) (**Table KB-1, Figures KB-1-3**).

Lower Kissimmee

For the week ending December 1, 2024, mean weekly discharge was 1,500 cfs and 1,400 cfs S-65 and S-65A, respectively. Mean weekly discharge from the Kissimmee River was 1,600 cfs and 1,300 cfs at S-65D and S-65E, respectively (**Table KB-2**). Mean weekly headwater stages were 45.2 feet NAVD88 at S-65A and 24.7 feet NAVD88 at S-65D. Mean weekly river channel stage decreased by 0.7 feet to 35.5 feet NAVD88 (**Figure KB-4**). Mean weekly water depth on the Kissimmee River floodplain decreased by 0.04 feet to 0.74 feet (**Table KB-2, Figure KB-5**). The weekly average concentration of dissolved oxygen in the Kissimmee River increased from 4.6 mg/L the previous week to 6.2 mg/L (**Table KB-2, Figure KB-6**).

Water Management Recommendations

Follow the Headwaters Revitalization Schedule (HRS) Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A (**Figure KB-7**). Maintain at least minimum flow (250-300 cfs) at S-65A.

Table KB-1. Average discharge for the preceding seven days, Sunday’s average daily stage and Sunday’s average daily departure from Kissimmee Chain of Lakes (KCOL) flood regulation lines or temporary schedules. All data are provisional.

Water Body	Structure	Stage Monitoring Site	Weekly (7-Day) Average Discharge (cfs)	Sunday Lake Stage (feet NAVD88) ^a	Schedule Type ^b	Sunday Schedule Stage (feet NAVD88)	Sunday Departure from Regulation (feet)	
							12/1/24	11/24/24
Lakes Hart and Mary Jane	S-62	LKMJ	0	60.0	R	59.9	0.1	0.0
Lakes Myrtle, Preston and Joel	S-57	S-57	8	61.0	R	61.0	0.0	-0.1
Alligator Chain	S-60	ALLI	0	62.9	R	63.0	-0.1	-0.1
Lake Gentry	S-63	LKGT	0	60.4	R	60.4	0.0	0.0
East Lake Toho	S-59	TOHOE	0	57.0	R	57.0	0.0	0.0
Lake Toho	S-61	TOHOW S-61	1	54.0	R	53.8	0.2	0.2
Lakes Kissimmee, Cypress and Hatchineha	S-65	KUB011 LKIS5B	1500	50.4	T	51.8	-1.4	-1.0

a. Names of in-lake monitoring sites and structures used to determine lake stage. If more than one site is listed, an average is reported.

b. A: projected recession line; R: USACE regulation schedule; S: temporary recession target line; T: temporary schedule; NA: not applicable or not available.




Figure KB-1. East Lake Toho regulation schedule, stage, discharge, and rainfall.




Figure KB-2. Lake Toho regulation schedule, stage, discharge, and rainfall.




Figure KB-3. Lakes Kissimmee, Cypress and Hatchineha regulation schedule, stage, discharge, and rainfall.

Table KB-2. One- and seven-day average discharge and stage at Lower Kissimmee basin structures, river channel dissolved oxygen concentrations and water depths in the Phase I area floodplain. All data are provisional.

Metric	Location	Sunday Daily Average	Weekly Average for Previous Seven Day Periods			
		12/1/24	12/1/24	11/24/24	11/17/24	11/10/24
Discharge	S-65	1,500	1,500	1,400	1,400	1,200
Discharge	S-65A ^a	1,400	1,400	1,400	1,400	1,300
Headwater Stage (feet NAVD88)	S-65A	45.3	45.2	45.2	45.2	45.2
Discharge	S-65D ^b	1,700	1,600	1,500	1,700	2,100
Headwater Stage (feet NAVD88)	S-65D ^c	24.7	24.7	25.0	25.6	26.5
Discharge (cfs)	S-65E ^d	1,300	1,300	1,500	1,800	2,200
Discharge (cfs)	S-67	0	0	0	0	0
Dissolved Oxygen (mg/L) ^e	Phase I, II/III river channel	6.4	6.2	4.6	3.7	3.2
River channel mean stage (feet NAVD88) ^f	Phase I river channel	35.5	35.5	36.2	36.0	36.5
Mean depth (feet) ^g	Phase I floodplain	0.73	0.74	0.78	0.86	1.10

a. Combined discharge from main and auxiliary structures.

b. Combined discharge from S-65D, S-65DX1, and S-65DX2.

c. Average stage from S-65D and S-65DX1.

d. Combined discharge from S-65E and S-65EX1.

e. Dissolved oxygen is the average of values from sondes KRBN, PC62, PC33, PD62R, and PD42R.

f. Mean of five river channel stations (PC62, KRDR02, KRBN, PC33, PC11) in the Phase I area.

g. One-day spatial average obtained from the South Florida Water Depth Assessment Tool (SFWDAT).




Figure KB-4. Kissimmee River stage, discharge, and rainfall.





Figure KB-5. Phase I-II-III area Kissimmee River floodplain water depths (from left to right) one year ago, one month ago, and current.



Report Date: 12/3/2024; data are through: 12/1/2024



Figure KB-6. Kissimmee River channel mean daily dissolved oxygen concentration (mg/L), S-65A discharge (cfs), temperature (°C) and rainfall (inches). Dissolved oxygen (DO) and temperature are mean daily values averaged for PC62, KRDR02, KRBN, PC33, PC11, PD62R, and PD42R with an average of five stations reporting this week. Rainfall values are daily totals for Kissimmee River (Pool BCD) AHED watershed.

HRS Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A

Discharge Guidance for Increment 1 Temporary Deviation Discharge Plan		
ZONE	S-65 RELEASES	S-65A TARGET FLOWS
ZONE A	Releases for Flood Risk Management up to maximum structure capacity as determined by downstream constraints with a firm capacity of 3,000 cfs.	
ZONE B1	1,400 cfs minimum ramp to 3,000 cfs at Zone A boundary	S-65A releases between 1,400 cfs and 3,000 cfs at Zone A boundary based on Table 1
ZONE B2	Releases as needed to target flows at S-65A	Target S-65A flows of 1,400 cfs to meet ecological needs
ZONE B3	Releases as needed to target flows at S-65A	S-65A flows between 300 cfs and 1,400 cfs
ZONE B4	Releases as needed to target flows at S-65A	Target S-65A flows of 300 cfs
ZONE B5	Releases as needed to target flows at S-65A	Target S-65A flows of 150 cfs
ZONE C	0 cfs	Flow as needed to maintain optimum S-65A headwater




Table KB-3. Maximum Rate of Change Limits for S-65A

MAXIMUM Release Rate of Change Limits for S-65A. In general recommended rates of change will be slower than shown in this table.

Q (cfs)	Maximum rate of INCREASE (cfs/day)	Maximum rate of DECREASE (cfs/day)
0-300	50	-50
301-650	75	-75
651-1400	150	-150
1401-3000	300	-600
>3000	1000	-2000

CENTRAL AND SOUTHERN FLORIDA PROJECT
 2024 LAKE KISSIMMEE, HATCHINEHA & CYPRESS
 Temporary Deviation (Increment 1)
 DATED: May 2024
 DEPARTMENT OF THE ARMY, JACKSONVILLE DISTRICT
 CORPS OF ENGINEERS, JACKSONVILLE, FLORIDA

Other Considerations

- When possible, limit lake ascension rate in the Jun 1 - Aug 15 window to 0.25 ft per 7 days in Lakes Kissimmee, Cypress, Hatchineha (S-65), East Toho (S-59) and Toho (S-61).
- If outlook is for extreme dry conditions meet with KB staff to discuss modifications to this plan.

Figure KB-7. Headwaters Revitalization Schedule (HRS) Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A.

Lake Okeechobee

Lake Okeechobee stage was 14.45 feet NAVD88 (15.76 ft NGVD29) on December 1, 2024, which was 0.15 feet lower than the previous week and 0.34 feet lower than a month ago (**Figure LO-1**). Lake stage is in Zone D of the regulation schedule (**Figure LO-2**) and was 0.76 feet above the upper limit of the recovery ecological envelope (**Figure LO-3**). According to NEXRAD, 0.0 inches of rain fell directly over the Lake last week.

Average daily inflows (excluding rainfall) decreased from 1,700 cfs the previous week, to 1,420 cfs. The largest single inflow came from the Kissimmee River via the S-65E structure (1,330 cfs). Average daily outflows (excluding evapotranspiration) increased from 2,550 cfs the previous week to 3,670 cfs. **Figures LO-4 and LO-5** show the combined average daily inflows and outflows for the Lake over the past eight weeks, and average inflows and outflows last week, respectively.

In the most recent non-obscured satellite image from November 28, 2024, NOAA's Harmful Algal Bloom Monitoring System suggests minimal bloom activity on Lake Okeechobee (**Figure LO-6**).

Note: All data presented in this report are provisional and are subject to change.

1 Month Ago:
11/01/2024

Current:
12/01/2024




Figure LO-1. Lake Okeechobee water depth estimates based on South Florida Water Depth Assessment Tool (SFWDAT).

Lake Okeechobee Water Level History and Projected Stages




Figure LO-2. Recent Lake Okeechobee stages with projected stages based on a dynamic position analysis. Note: stages are in NGVD29, approximate NAVD88 values are shown for reference.

Lake Okeechobee Stage vs Recovery Ecological Envelope




Figure LO-3. The current and seven prior year's annual stage hydrographs for Lake Okeechobee in comparison to the recovery envelope (light green). A shift from the normal ecological envelope to the recovery envelope occurred because the 30-day minimum lake stage (elevations exposed for at least 30 days, nonconsecutively) in the June 1 – July 31, 2023, window was >13 ft NGVD29 (11.75 ft NAVD88).




Figure LO-4. Major inflows (orange) to and outflows east and west (blue) from Lake Okeechobee. Outflows south are shown in green. Flows into Lake Okeechobee from the L-8 canal through S-271 (formerly Culvert 10A) or from the C-44 canal through the S-308 are included as inflows. Conversely, flows from Lake Okeechobee into the L-8 or C-44 canals are included with outflows. Inflows are shown as positive values; outflows are negative. Outflows through the S-77 (Caloosahatchee) and S-308 (C-44 Canal) structures are based on downstream gauges to include flows to lock openings for navigation.




Figure LO-5. Inflows into Lake Okeechobee from Indian Prairie basins, Taylor Creek/Nubbin Slough, Kissimmee River and Fisheating Creek, and outflows to the west via S-77, to the east via S-308, to the south via S-351, S-352, S-354, and to southeast via S-271 (formerly Culvert 10A) for the week of November 25 – December 01, 2024.




Figure LO-6. Cyanobacteria bloom index level on Lake Okeechobee, based on NOAA’s harmful algal bloom monitoring system. Gray color indicates cloud cover. *Provisional NOAA image, subject to change*.

Estuaries

St. Lucie Estuary

Mean total inflow to the St. Lucie Estuary was 289 cfs from November 26 to December 1, 2024 (**Figures ES-1 and ES-2**), and the week prior inflow was 346 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-2**.

Over the past two weeks, salinities generally increased at all three sites within the estuary (**Table ES-1 and Figure ES-3**). The seven-day moving average of the surface and bottom salinities at the US1 Bridge was 18.1. Salinity conditions in the middle estuary were estimated to be within the optimal range for adult eastern oysters (**Figure ES-4**). The mean larval oyster recruitment rate reported by the Fish and Wildlife Research Institute (FWRI) was 0.4 spat/shell for October, which was similar to the previous month (**Figure ES-5**).

Caloosahatchee River Estuary

Mean total inflow to the Caloosahatchee River Estuary was 2,083 cfs from November 26 to December 1, 2024 (**Figures ES-6 and ES-7**), and the week prior total inflow was 2,168 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-7**.

Over the past two weeks, salinities continued to decrease at all sites within the estuary (**Table ES-2 and Figures ES-8 and ES-9**). Fort Myers surface sensor is currently offline and not reporting this week. The seven-day mean salinities (**Table ES-2**) were in the optimal range (0-10) for tape grass in the upper estuary. The seven-day mean salinity values were within the optimal range for adult eastern oysters at Cape Coral and Shell Point and in the upper stressed range at Sanibel (**Figure ES-10**). The mean larval oyster recruitment rate reported by the FWRI was 1.8 spat/shell at Iona Cove and 16.6 spat/shell at Bird Island for October, which was a decrease at both sites from the previous month, indicating the end of fall spawning season (**Figures ES-11 and ES-12**).

Surface salinity at Val I-75 was forecast for the next two weeks using an autoregression model (Qiu and Wan, 2013¹) coupled with a linear reservoir model for the tidal basin. Model scenarios included pulse releases at S-79 ranging from 450 to 2,000 cfs with estimated tidal basin inflows of 117 cfs. Model results from all scenarios predict daily salinity to be 0.5 or lower and the 30-day moving average surface salinity to be 0.8 or lower at Val I-75 at the end of the two-week period (**Table ES-3 and Figure ES-13**). This keeps predicted salinities in the upper estuary within the optimal salinity range (0-10) for tape grass.

¹ Qiu, C., and Y. Wan. 2013. Time series modeling and prediction of salinity in the Caloosahatchee River Estuary. *Water Resources Research* 49:5804-5816.

Red Tide

The FWRI reported on November 26, 2024, that *Karenia brevis*, the Florida red tide dinoflagellate, was observed at low concentrations in samples collected from Charlotte County over the past week and was not present in any other samples collected within the District region.

Water Management Recommendations

Lake stage is in Zone D. Current climatological and hydrological conditions are normal. The LOSOM release guidance suggests up to 2,000 cfs release at S-79 to the Caloosahatchee River Estuary and no releases at S-80 to the St. Lucie Estuary.




Figure ES-1. Basins, water control structures and salinity monitoring sites in the St. Lucie Estuary.




Figure ES-2. Total daily inflows from Lake Okeechobee and runoff from the C-44, C-23, C-24, Ten Mile Creek, and Tidal Basins into the St. Lucie Estuary.

Table ES-1. Seven-day mean salinity at oyster monitoring sites in the St. Lucie Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope reflects the optimum salinity range for adult eastern oysters (*Crassostrea virginica*) in the estuary. Data are provisional.

Sampling Site	Surface	Bottom	Optimum Envelope
HR1 (North Fork)	13.7 (9.0)	16.4 (13.6)	10.0 – 25.0
US1 Bridge	17.1 (14.0)	19.1 (16.9)	10.0 – 25.0
A1A Bridge	24.4 (21.7)	27.1 (26.3)	10.0 – 25.0




Figure ES-3. Mean daily salinity at the A1A, US1, and HR1 sites in the St. Lucie Estuary.




Figure ES-4. Seven-day moving average of the surface and bottom salinities at the US1 Bridge in the St. Lucie Estuary.




Figure ES-5. Mean oyster recruitment at the Rio oyster monitoring station and 14-day mean salinity at US1 Bridge.




Figure ES-6. Basins, water control structures, and salinity monitoring sites in the Caloosahatchee River Estuary.




Figure ES-7. Total daily inflows from Lake Okeechobee, and runoff from the C-43 and Tidal basins into the Caloosahatchee River Estuary.

Table ES-2. Seven-day mean salinity at six monitoring sites in the Caloosahatchee River Estuary. Current means are in bold font; previous week's means are in parentheses. Missing or not reported values are represented by NR. The envelope in the upper estuary sites is for the protection of tape grass and the envelope in the lower estuary is the optimum salinity range for adult eastern oysters (*Crassostrea virginica*). Data are provisional.

Sampling Site	Surface	Bottom	Optimum Envelope
S-79 (Franklin Lock)	0.7 (0.8)	0.7 (0.8)	0.0 – 10.0
Val I-75	1.0 (1.3)	1.6 (2.2)	0.0 – 10.0
Fort Myers Yacht Basin	NR (NR)	7.6 (11.9)	0.0 – 10.0
Cape Coral	10.7 (14.0)	12.7 (16.6)	10.0 – 25.0
Shell Point	22.8 (25.5)	23.6 (26.6)	10.0 – 25.0
Sanibel	28.0 (29.8)	28.9 (31.1)	10.0 – 25.0




Figure ES-8. Mean daily salinity at upper Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




Figure ES-9. Mean daily surface salinity at lower Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




Figure ES-10. Seven-day moving average of surface and bottom salinities at Cape Coral, Shell Point, and Sanibel monitoring sites in the Caloosahatchee River Estuary.




Figure ES-11. Mean oyster recruitment at the Iona Cove oyster monitoring station and 14-day mean salinity at Cape Coral.




Figure ES-12. Mean oyster recruitment at the Bird Island oyster monitoring station and 14-day mean salinity at Shell Point.

Table ES-3. Predicted salinity at Val I-75 in the Caloosahatchee River Estuary at the end of the forecast period for various S-79 flow release scenarios.

Scenario	Simulated S-79 Flow (cfs)	Tidal Basin Runoff (cfs)	Daily Salinity	30-Day Mean Salinity
A	450	117	0.5	0.8
B	650	117	0.4	0.8
C	1,200	117	0.3	0.8
D	2,000	117	0.3	0.8

Observed and Forecasted Flow at S-79 and Salinity at Val I-75




Figure ES-13. Surface salinity forecast at the Val I-75 site assuming a 450 cfs pulse release at S-79.

Stormwater Treatment Areas

STA-1E: STA-1E Central Flow-way is offline for construction activities. An operational restriction is in place in the Western Flow-way for post-construction vegetation grow-in. Online treatment cells are above target stage. (**Figure S-1**).

STA-1W: An operational restriction is in place in the Northern Flow-way for vegetation management activities. Treatment cells are above target stage. Vegetation in the flow-ways is highly stressed. The 365-day PLRs for the Eastern and Western Flow-ways are high, and the 365-day PLR for the Northern Flow-ways is below $1.0 \text{ g/m}^2/\text{year}$ (**Figure S-1**).


STA-2: Operational restrictions are in place in Flow-ways 2 and 4 for vegetation management activities. Online treatment cells are above target stage. Vegetation in Flow-ways 2, 3, and 4 is stressed, and in 5 is highly stressed. The 365-day PLRs for Flow-ways 1, 4, and 5 are below $1.0 \text{ g/m}^2/\text{year}$. The 365-day PLRs for Flow-ways 2 and 3 are high (**Figure S-2**).

STA-3/4: An operational restriction is in place in the Eastern Flow-way for post-drawdown vegetation grow-in. Treatment cells are above target stage. Vegetation in the Central Flow-way is highly stressed and in the Eastern Flow-way is stressed. The 365-day PLRs for the Central and Western Flow-ways are high (**Figure S-2**).

STA-5/6: Treatment cells are above target stage. All treatment cells have highly stressed or stressed vegetation conditions. The 365-day PLRs for Flow-ways 1, 6, 7, and 8 are below $1.0 \text{ g/m}^2/\text{year}$, and the 365-day PLRs for Flow-ways 2, 3, 4, and 5 are high. (**Figure S-3**).

For definitions on STA operational language see glossary following figures.

Eastern Flow Path Weekly Status Report – 11/25/2024 through 12/1/2024




STA-1W	Flow-way Status
Western	<ul style="list-style-type: none"> • High 365-day PLR • Highly stressed vegetation conditions
Eastern	<ul style="list-style-type: none"> • High 365-day PLR • Highly stressed vegetation conditions
Northern	<ul style="list-style-type: none"> • Stressed vegetation conditions • Planting emergent vegetation
Cell 6	
Cell 7+8	

STA-1E	Flow-way Status
Western	<ul style="list-style-type: none"> • Post-construction vegetation grow-in
Central	<ul style="list-style-type: none"> • Offline for construction activities
Eastern	

Figure S-1. Eastern Flow Path Weekly Status Report

Central Flow Path Weekly Status Report – 11/25/2024 through 12/1/2024



STA-3/4	Flow-way Status
Western	<ul style="list-style-type: none"> High 365-day PLR
Central	<ul style="list-style-type: none"> Highly stressed vegetation conditions High 365-day PLR
Eastern	<ul style="list-style-type: none"> Post-drawdown vegetation grow-in

STA-2	Flow-way Status
Flow-way 1	<ul style="list-style-type: none"> Upstream FAV control
Flow-way 2	<ul style="list-style-type: none"> High 365-day PLR Post-construction vegetation grow-in Stressed vegetation conditions Upstream FAV control
Flow-way 3	<ul style="list-style-type: none"> High 365-day PLR Stressed vegetation conditions Upstream FAV control
Flow-way 4	<ul style="list-style-type: none"> Planting emergent vegetation Upstream FAV control
Flow-way 5	<ul style="list-style-type: none"> Highly stressed vegetation conditions

Figure S-2. Central Flow Path Weekly Status Report

Western Flow Path Weekly Status Report – 11/25/2024 through 12/1/2024




Figure S-3. Western Flow Path Weekly Status Report

Basic Concepts and Definitions for STA Weekly Status Report

- **Inflow:** Sum of flow volume at all inflow structures to an STA.
- **Lake Inflow:** Portion of the STA total inflow volume that originates from Lake Okeechobee.
- **Outflow:** Sum of flow volume at outflow structures from an STA.
- **Total Phosphorus (TP):** Total mass of phosphorus in all its forms; including particulate, dissolved, etc.
- **Inflow Concentration:** TP concentration is the mass of TP in micrograms per liter of water, $\mu\text{g/L}$ or ppb. Inflow concentration refers to the flow-weighted mean TP from all inflow structures over a period of time.
- **Outflow Concentration:** The flow-weighted mean TP from all outflow structures over a period of time. The outflow concentration represents the reduction of inflow TP achieved by STA treatment of the inflow water.
- **WQBEL:** The STA outflow concentration that is required upon completion of the Restoration Strategies projects by December 2025. The outflow concentration shall not exceed 13 ppb as an annual flow weighted mean in more than 3 out of 5 water years on a rolling basis and shall not exceed 19 ppb as an annual flow weighted in any water year.
- **Flow-Way (FW):** One or more treatment cells connected in series. Cells typically have emergent aquatic vegetation (EAV) in the front portion of the flow-way followed by a mix of EAV and submerged aquatic vegetation (SAV)
- **Vegetation Status:** Healthy means the vegetation condition is good and will allow the STA to perform as designed. Stressed means the vegetation is showing signs of poor health, such as browning or areas of vegetation die-off, or the cell contains undesirable vegetation such as floating exotic vegetation requiring treatment. The TP reduction capability of the STA is affected when the vegetation condition is poor.
- **Phosphorus Loading Rate (PLR):** Mass of inflow TP in grams, divided by total treatment area of STA in square meters, per year. In general, a 365-day value of less than 1.0 is needed for an STA to perform optimally. A PLR of 2.0 is considered very high and a PLR of 3.0 is considered extremely high. The TP reduction capability of the STA is affected when the PLR is high, very high and extremely high.
- **Online:** Online status means the FW can receive and treat inflow.
- **Online with Restriction:** The FW can receive and treat inflow, but the amount of flow or water level may be limited temporarily. For example, a vegetation rehabilitation effort may require reduced flows through an area while the new plants are establishing, or nesting by protected species may require a certain water level not to be exceeded.
- **Offline:** The FW is unable to receive and treat inflow due to repairs, construction, or other prohibitive reasons.
- **Depth:** Difference between the average surface water level in a cell and the average ground elevation in that cell. Target depths, or depths between flow events, are between 1.25 ft to 1.5 ft. As depth approaches or drops below zero, an increasing percentage of the cell is considered dry and STA conditions deteriorate. An increase in depth above target depth is expected with increasing flow. However, as depth increases much above the target depth and is sustained over a period of time, it can be detrimental to vegetation health and overall STA treatment performance.
- **Note:** The data provided in this summary report were developed using a combination of provisional and quality-assured flow and water quality data. In some cases, best professional judgment was used to estimate missing data and revise questionable data. Values provided are not considered final but are appropriate for use in STA operational decision-making.

Everglades

Water Conservation Area Regulation Schedules

No rainfall occurred throughout the Everglades Protection Area (EPA). WCA-1: Stages within the Refuge continue to decline and remains below the A2 zone regulation line. On Sunday, the 3-Gauge average was 0.25 feet below the A2 Zone regulation line. WCA-2A: Stage ascension at Gauge 2A-17 remains above but following the slope of the regulation line. The average on Sunday was around 1.0 feet above the Zone A line. WCA-3A: The 3-Gauge average stage continued to decline over the last week, now falling just below the Zone A regulation line by around 0.1 feet on Sunday. WCA-3A North: Stage at Gauge 62 (NW corner) continues to recede away from the regulation line last week maintaining a declining slope. The average on Sunday was 0.56 feet below the Upper Schedule. See figures **EV-1** through **EV-4**.

Water Depths

The SFWDAT model output for December 1, 2024, illustrates shallower conditions throughout the EPA. Notably, there is a continued expansion of drying conditions in the northern portions of the WCAs and a contraction of ponded conditions in southern/eastern WCA3A and in northern Shark River Slough (SRS). Big Cypress has greater potential for water below ground surface which is expanding both to the north and south of Tamiami trail. Hydrologic connectivity has shown a reduction in Shark River Slough, and Taylor Slough has shown a potential closure of flow as depths are falling below ground along the path of the slough. Current WDAT water depth estimates when compared to one month ago, indicate a uniform pattern of shallower conditions throughout the EPA with greatest recessions occurring from WCA2A into the entirety of WCA3A and out west into Big Cypress area. Southern Everglades National Park (ENP) is further experiencing recessions of nearly a foot over the past month, particularly in the vicinity of Taylor Slough. The comparison to modeled conditions a year ago show similar shallower conditions throughout the EPA. Comparing current conditions to the 20-year percentiles for December 1st, depth conditions moved towards average in WCA-1, below average in northern WCA-2A, and the majority of WCA-3A has fallen below the 70th percentile. Most of ENP and WCA-3B remain above the 80th percentile. Southern ENP in Taylor Slough now falls below average. See figures **EV-5** through **EV-7**.

Taylor Slough and Florida Bay

All stages decreased across Taylor Slough over the past week, with an average decrease of 0.20 feet. Changes ranged from -0.41 feet at Taylor Slough Bridge (TSB) in the northern slough to -0.08 feet EPSW in the C-111 area (**Figure EV-8** and **Figure EV-9**). Taylor Slough water levels remain above the recent average for this time of year by 3.1 inches compared to before the Florida Bay initiative (starting in 2017). The Craighead Pond (CP) and TSB stages are below estimated historical levels (circa 1900) by 0.44 and 1.21 feet, respectively.

Average Florida Bay salinity was 21.5. Salinity increased at most sites, with changes ranging from -6.1 at Garfield Bight (GB) in the western nearshore region to +9.7 at Buoy Key (BK) in the western offshore region (**Figure EV-8**). Salinity is within the WY2001-2016 interquartile range (IQR) in the eastern region but remains below the IQR and above

estimated historical levels (circa 1900) in the central and western regions (**Figure EV-10**). Average Florida Bay salinity remains below its recent average for this time of year by 2.6.

Salinity at the Taylor River (TR) station in the mangrove zone (tracked for the Florida Bay MFL) was 0.3. The 30-day moving average was 0.5 (**Figure EV-11**). The 365-day moving sum of flow from the five major creeks (McCormick Creek, Taylor River, Mud Creek, Trout Creek, West Highway Creek) was 406,091 acre-feet (**Figure EV-11**).

Average rainfall across Taylor Slough and Florida Bay was 0.0 inches over the past week, based on the 18 gauges used for this report (**Figure EV-12**). Wind directions and speeds in Florida Bay ranged from 3.6 mph W on November 29th to 21.7 mph N on November 30th (**Figure EV-12**).

Average daily flow from the five major creeks totaled 1,498 acre-feet last week, with net positive flows for the week. Total daily creek flow ranged from 195 acre-feet on November 28th to 2,499 acre-feet on December 1st (**Figure EV-13**). Average daily flow for the week was 2,499 acre-feet below estimated historical levels (circa 1900).

Implications for water management

The ecology of the Everglades benefits from recession rates from 0.0 to 0.12 feet per week this time of year. Maintaining a hydroperiod supportive of upcoming wading bird nesting at the Alley North colony in WCA-3A North is less likely as drier conditions persist in that region as we approach La Nina climatic conditions. Florida Bay salinity is at a good position as we continue into the dry season and will continue to benefit from maintaining freshwater input to the system when available. Maintaining higher water depths in the WCAs early in the dry season can help support wading bird nesting success. Although reducing water flow to ENP could lead to increased salinity in the bay, current favorable salinity levels may provide an opportunity for a temporary reduction in inflow from the WCAs to the park. This approach could deliver broader landscape-wide ecological benefits across the EPA while balancing ecological priorities. Individual regional recommendations can be found in **Table EV-2**.

Table EV-2. Previous week's rainfall and water depth changes in Everglades basins.

Everglades Region	Rainfall (inches)	Stage change (feet)
WCA-1	0.00	-0.04
WCA-2A	0.00	-0.11
WCA-2B	0.00	-0.12
WCA-3A	0.00	-0.16
WCA-3B	0.00	-0.09
ENP	0.00	-0.01




Figure EV-1. WCA-1 stage hydrographs and regulation schedule.




Figure EV-2. WCA-2A stage hydrographs and regulation schedule.




Figure EV-3. WCA-3A stage hydrographs (three-gauge average, 3-69W) and regulation schedule.




Figure EV-4. WCA-3A stage hydrograph (Deer gauge; Site 62) and GA62 regulation schedule.




Figure EV-5. Everglades water depths from two months ago (left), one month ago (center) and present (right), based on SFWDAT.




Figure EV-6. Present Everglades water depths (left) and water depth changes from one month (center) and one year (right) ago, based on SFWDAT.




Figure EV-7. Present water depths (November 10th, 2024) compared to the day of year average over the previous 20 years.




Figure EV-8. Taylor Slough water depths and Florida Bay salinities with changes from a week ago.




Figure EV-9. Taylor Slough water depth time series for Taylor Slough Bridge (TSB; northern slough) and Craighhead Pond (CP; southern slough).




Figure EV-10. Eastern (top panel), Central (middle panel) and Western (bottom panel) Florida Bay daily average salinities with WY2001-2016 interquartile (25-75 percentile) ranges (IQR) and estimated historical daily average salinities. The hypersalinity threshold indicates the level at which salinities start to become harmful to seagrass.




Figure EV-11. Salinity at Taylor River (TR; top) and creek inflow to Florida Bay (bottom) from the five major creeks (McCormick Creek, Taylor River, Mud Creek, Trout Creek, and West Highway Creek). The 30-day moving average salinity and 365-day total creek flow are tracked for the Florida Bay MFL criteria.




Figure EV-12. Daily average rain across Taylor Slough and Florida Bay, along with hourly average wind speed and direction (measured at Long Key) in Florida Bay over the past four weeks.




Figure EV-13. Daily average creek flow summed between five creeks with estimated historical daily flow over the past four weeks.

Table EV-2. Weekly water depth changes and water management recommendations

SFWMD Everglades Ecological Recommendations, December 2, 2024 (red is new)			
	Weekly change	Recommendation	Reasons
WCA-1	Stage decreased by 0.04 feet	Recession rate of less than 0.06 feet per week.	Protect within basin and downstream habitat and wildlife.
WCA-2A	Stage decreased by 0.11 feet	Recession rate of less than 0.12 feet per week.	Protect within basin and downstream habitat and wildlife.
WCA-2B	Stage decreased by 0.12 feet	Recession rate of less than 0.12 feet per week.	Protect within basin and downstream habitat and wildlife.
WCA-3A NE	Stage decreased by 0.20 feet	Recession rate of less than 0.12 feet per week.	Protect within basin and downstream habitat and wildlife.
WCA-3A NW	Stage decreased by 0.13 feet	Recession rate of less than 0.12 feet per week.	
Central WCA-3A S	Stage decreased by 0.17 feet	Recession rate of less than 0.12 feet per week.	Protect within basin wildlife.
Southern WCA-3A S	Stage decreased by 0.12 feet		
WCA-3B	Stage decreased by 0.09 feet	Recession rate of less than 0.12 feet per week.	Protect within basin and downstream habitat and wildlife.
ENP-SRS	Stage decreased by 0.01 feet	Make discharges to ENP according to COP and TTFF protocol while adaptively considering upstream and downstream ecological conditions.	Protect within basin and upstream habitat and wildlife.
Taylor Slough	Stage changes ranged from -0.41 feet to -0.08 feet	Move water southward as possible.	When available, provide freshwater to promote water movement.
FB- Salinity	Salinity changes ranged from -6.1 to +9.7	Move water southward as possible.	When available, provide freshwater to promote water movement.

Biscayne Bay

As shown in **Figure BB-1**, mean total inflow to Biscayne Bay was 299 cfs, and the previous 30-day mean inflow was 338 cfs. The seven-day mean salinity was 28.9 at BBCW8 and 23.6 at BBCW10, both within the ideal salinity range for estuarine organisms in this region (salinity less than 35). Data were provided by Biscayne National Park.




Figure BB-1. Seven-day mean salinity at BBCW8 and BBCW10 and total daily flow in Biscayne Bay. Total daily flow was calculated using flow from structures S20G, S20F, S21, S21A, S123, and S700P.