

Disclaimer: Information contained in the report addresses environmental conditions only and is not the official South Florida Water Management District operations recommendation or decision.

## **M E M O R A N D U M**

**TO:** John Mitnik, Assistant Executive Director, Executive Office Staff

**FROM:** SFWMD Staff Environmental Advisory Team

**DATE:** January 28, 2026

**SUBJECT:** Weekly Environmental Conditions for Systems Operations

### **Summary**

#### **Weather Conditions and Forecast**

Atmospheric disturbance systems will dig southeastward across Florida on Wednesday night. As it approaches, moisture levels may recover slightly across the southeastern portion of Florida, potentially supporting the development of generally light, isolated showers along the lower east coast and the Florida Keys on Wednesday into Thursday. Elsewhere, dry conditions will prevail. Fast-moving light showers are expected to reach the Kissimmee Valley late Friday night, then progress into the Lake Okeechobee region by Saturday morning before clearing the lower east coast by Saturday afternoon. Due to the fast speed of these showers, significant rainfall is unlikely. A strong cold front will surge southward into Florida Saturday afternoon. The frontal passage is expected to be accompanied by very gusty northwesterly winds of 20–25 mph, with gusts potentially reaching 35–40 mph. A very cold and dry Arctic air mass will follow, leading to steadily falling temperatures throughout the day with high rates of evapotranspiration. There is a growing signal that freezing temperatures could reach as far south as the Miami area Saturday night and again Sunday night. Following the frontal passage, the wind will gradually relax by Monday. Cool and dry conditions are expected to persist across most of Florida into Monday with no rainfall expected. For the 7-day period ending Tuesday morning, much below average total SFWMD weekly rainfall is expected.

#### **Kissimmee**

In the past week, releases were made as needed from East Lake Toho and Lake Toho to continue snail kite nesting season stage recessions to reach low pool by June 1, 2026. Releases from KCH followed the Headwaters Revitalization Schedule (HRS) Increment 1 Temporary Deviation Discharge Plan. Weekly average discharge on January 25, 2026, was 450 cfs at S-65 and 390 cfs at S-65A, respectively. Mean weekly water depth on the Kissimmee River floodplain decreased by 0.01 feet to 0.35 feet. The weekly average concentration of dissolved oxygen in the Kissimmee River increased from the previous week's value of 8.2 mg/L to 8.6 mg/L, which is above both the potentially lethal level of 1.0 mg/L and the stressful level of 2.0 mg/L for Florida bass and other species.

## **Lake Okeechobee**

Lake Okeechobee stage was 11.68 feet NAVD88 (12.98 ft NGVD29) on January 25, 2026, which was 0.10 feet lower than the previous week and 0.49 feet lower than a month ago. Average daily inflows (excluding rainfall) decreased from the previous week, dropping from 400 cfs to 320 cfs. Average daily outflows (excluding evapotranspiration) increased slightly from the previous week, going from 2,020 cfs to 2,090 cfs. The most recent non-obscured satellite image from January 25, 2026, NOAA's Harmful Algal Bloom Monitoring System suggests moderate to high cyanobacteria activity in the southern and western regions of the lake. The January 10-11 phytoplankton sampling showed 2 of 8 samples had detectable levels of microcystins, though both were < 0.5 µg/L, and 5 of the 30 water quality samples had chlorophyll a > 40 µg/L, 3 of which had values > 80 µg/L (all from the south end of the lake).

## **Estuaries**

Total inflow to the St. Lucie Estuary averaged 110 cfs over the past week with no flow coming from Lake Okeechobee. Mean salinities increased at all sites over the past week. Salinity in the middle estuary was in the upper stressed range (> 25) for adult eastern oysters.

Total inflow to the Caloosahatchee Estuary averaged 458 cfs over the past week with 249 cfs coming from Lake Okeechobee. Over the past week, salinities increased at Shell Point and Sanibel and decreased at the remaining sites. Salinities were in the optimal range (0-10) for tape grass at S-79 and Val I-75 and in the stressed range (10-15) at Fort Myers. Salinities were in the optimal range (10-25) for adult oysters at Cape Coral and in the upper stressed ranged (>25) at Shell Point and Sanibel.

## **Stormwater Treatment Areas**

For the week ending Sunday, January 25<sup>th</sup>, 2026, 6,500 ac ft of Lake Okeechobee water was delivered to the FEBs/STAs. The total amount of Lake releases sent to the FEBs/STAs in WY2026 is approximately 48,100 ac-feet. The total amount of inflows to the STAs in WY2026 is approximately 551,000 ac-feet. Online STA treatment cells are at or above target stage. STA-1E Central Flow-way is offline for construction activities. STA-1W Eastern Flow-way is offline for vegetation management activities. Operational restrictions are in effect in STA-1E Western Flow-way, STA-2 Flow-ways 2, 3 and 4, and STA-3/4 Eastern Flow-way for vegetation management activities. This week, if LOSOM recommends Lake releases to the WCAs and conditions allow, releases will be sent to STA-2, STA-3/4 and STA-5/6.

## **Everglades**

Over the past week, the Everglades Protection Area (EPA) primarily received below average rainfall with exception of two days that were above average. Water depth recessions increased across the EPA. Recession rates at monitored sites increased last week to an average of approximately -0.09 ft/week, hindering water conservation during a period of below-average water depths across the Everglades. Southern WCA-2A remains unseasonably deep, while most of WCA-3A and ENP continue to be dry, below the 10th percentile, as it has for much of the water year. Below-average depths in WCA-3A and ENP limit aquatic prey production, increase the risk of damaging wildfire, promote

peat oxidation, and ridge/slough degradation. Wading bird activity remains limited, with foraging concentrated along the SW ENP coast and western Florida Bay during very low tides, while most major colonies across ENP and the WCAs remain inactive. Localized foraging by white ibis and spoonbills is occurring near Alley N, where approximately 100 pairs of spoonbills have begun nest platform construction, marking the onset of the nesting season. Taylor Slough stages continued to drop last week and remain well below the recent averages for this time of year. Average Florida Bay salinities increased last week and remains above recent averages for this time of year.

## Supporting Information

### Kissimmee Basin

#### *Upper Kissimmee*

On January 25, 2026, mean daily lake stages were 56.5 feet NAVD88 (0.5 feet below schedule) in East Lake Toho, 53.6 feet NAVD88 (0.4 feet below schedule) in Lake Toho, and 48.6 feet NAVD88 (3.0 feet below the Increment 1 Temporary Deviation schedule) in Lakes Kissimmee-Cypress-Hatchineha (KCH) (**Table KB-1, Figures KB-1-3**).

#### *Lower Kissimmee*

For the week ending January 25, 2026, mean weekly discharge was 450 cfs at S-65 and 390 cfs at S-65A. Mean weekly discharge from the Kissimmee River was 430 cfs at S-65D and 320 at S-65E (**Table KB-2**). Mean weekly headwater stages were 45.2 feet NAVD88 at S-65A and 29.1 feet NAVD88 at S-65D. Mean weekly river channel stage decreased by 0.4 feet from the previous week's value of 32.1 feet to 31.7 feet NAVD88 (**Figure KB-4**). Mean weekly water depth on the Kissimmee River floodplain decreased by 0.01 feet to 0.35 feet (**Table KB-2, Figure KB-5**). The weekly average concentration of dissolved oxygen in the Kissimmee River increased from 8.2 mg/L the previous week to 8.6 mg/L (**Table KB-2, Figure KB-6**).

#### **Water Management Recommendations**


Continue the stage recessions in East Lake Toho and Lake Toho lakes to reach their low pools on June 1, 2026. In KCH, follow the Headwaters Revitalization Schedule (HRS) Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A (**Figure KB-7**). With KCH stage near the bottom of Zone B3, target flows between 300 and 1,400 cfs at S-65A, using the Increment 1 Interpolation Tool to determine discharge relative to stage in KCH. If stage increases into Zone B2, target flows of 1,400 cfs; if stage decreases into Zone B4, target flows of 300 cfs.

**Table KB-1.** Average discharge for the preceding seven days, Sunday's average daily stage and Sunday's average daily departure from Kissimmee Chain of Lakes (KCOL) flood regulation lines or temporary schedules. All data are provisional.


Water Body	Structure	Stage Monitoring Site	Weekly (7-Day) Average Discharge (cfs)	Sunday Lake Stage (feet NAVD88) <sup>a</sup>	Schedule Type <sup>b</sup>	Sunday Schedule Stage (feet NAVD88)	Sunday Departure from Regulation (feet)	
							1/25/26	1/18/26
Lakes Hart and Mary Jane	S-62	LKMJ	0	60.1	R	59.9	0.2	0.1
Lakes Myrtle, Preston and Joel	S-57	S-57	11	60.3	R	60.3	0.0	0.0
Alligator Chain	S-60	ALLI	0	62.9	R	62.9	0.0	0.0
Lake Gentry	S-63	LKGT	0	60.4	R	60.4	0.0	0.0
East Lake Toho	S-59	TOHOE	140	56.5	R	57.0	-0.5	-0.3
Lake Toho	S-61	TOHOW S-61	370	53.6	R	54.0	-0.4	-0.3
Lakes Kissimmee, Cypress and Hatchineha	S-65	KUB011 LKIS5B	450	48.6	T	51.6	-3.0	-3.0

a. Names of in-lake monitoring sites and structures used to determine lake stage. If more than one site is listed, an average is reported.


b. A: projected recession line; R: USACE regulation schedule; S: temporary recession target line; T: temporary schedule; NA: not applicable or not available.



**Figure KB-1.** East Lake Toho regulation schedule, stage, discharge, and rainfall.



**Figure KB-2.** Lake Toho regulation schedule, stage, discharge, and rainfall.



**Figure KB-3.** Lakes Kissimmee, Cypress and Hatchineha regulation schedule, stage, discharge, and rainfall.

**Table KB-2.** One- and seven-day average discharge and stage at Lower Kissimmee basin structures, river channel dissolved oxygen concentrations and water depths in the Phase I area floodplain. All data are provisional.

Metric	Location	Sunday Daily Average	Weekly Average for Previous Seven Day Periods				
		1/25/26	1/25/26	1/18/26	1/11/26	1/4/26	
Discharge	S-65	420	450	530	500	420	
Discharge	S-65A <sup>a</sup>	370	390	470	430	370	
Headwater Stage (feet NAVD88)	S-65A	45.3	45.2	45.2	45.2	45.2	
Discharge	S-65D <sup>b</sup>	360	430	510	410	470	
Headwater Stage (feet NAVD88)	S-65D <sup>c</sup>	24.7	29.1	29.4	29.0	29.3	
Discharge (cfs)	S-65E <sup>d</sup>	300	320	400	290	370	
Discharge (cfs)	S-67	0	0	0	0	0	
Dissolved Oxygen (mg/L) <sup>e</sup>	Phase I, II/III river channel	8.4	8.6	8.2	7.8	8.5	
River channel mean stage (feet NAVD88) <sup>f</sup>	Phase I river channel	31.5	31.7	32.1	31.6	31.8	
Mean depth (feet) <sup>g</sup>	Phase I & II/III floodplain	0.35	0.35	0.36	0.35	0.36	

a. Combined discharge from main and auxiliary structures.

b. Combined discharge from S-65D, S-65DX1, and S-65DX2.

c. Average stage from S-65D and S-65DX1.


d. Combined discharge from S-65E and S-65EX1.

e. Dissolved oxygen is the average of values from sondes KRBn, PC62, PC33, PD62R, and PD42R.

f. Mean of five river channel stations (PC62, KRDR02, KRBn, PC33, PC11) in the Phase I area.


g. One-day spatial average obtained from the South Florida Water Depth Assessment Tool (SFWDAT).

### Stage and Discharge Hydrograph (report date: 1/27/2026; data through: 1/25/2026)




\*River Channel Stage is the average for PC62, KRDR02, KRBN, PC33, and PC11.

**Figure KB-4.** Kissimmee River stage, discharge, and rainfall.



**Figure KB-5.** Phase I-II-III area Kissimmee River floodplain water depths (from left to right) one year ago, one month ago, and current.



**Figure KB-6.** Kissimmee River channel mean daily dissolved oxygen concentration (mg/L), S-65A discharge (cfs), temperature (°C) and rainfall (inches). Dissolved oxygen (DO) and temperature are mean daily values averaged for PC62, KRDR02, KRBN, PC33, PC11, PD62R, and PD42R with an average of five stations reporting this week. Rainfall values are daily totals for Kissimmee River (Pool BCD) AHED watershed.

## HRS Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A

## Discharge Guidance for Increment I Temporary Deviation Discharge Plan


ZONE	S-65 RELEASES	S-65A TARGET FLOWS
ZONE A	Releases for Flood Risk Management up to maximum structure capacity as determined by downstream constraints with a firm capacity of 3,000 cfs.	
ZONE B1	1,400 cfs minimum ramp to 3,000 cfs at Zone A boundary	S-65A releases between 1,400 cfs and 3,000 cfs at Zone A boundary based on Table 1
ZONE B2	Releases as needed to target flows at S-65A	Target S-65A flows of 1,400 cfs to meet ecological needs
ZONE B3	Releases as needed to target flows at S-65A	S-65A flows between 300 cfs and 1,400 cfs
ZONE B4	Releases as needed to target flows at S-65A	Target S-65A flows of 300 cfs
ZONE B5	Releases as needed to target flows at S-65A	Target S-65A flows of 150 cfs
ZONE C	0 cfs	Flow as needed to maintain optimum S-65A headwater

Table KB-3. Maximum Rate of Change Limits for S-65A

MAXIMUM Release Rate of Change Limits for S-65A. In general recommended rates of change will be slower than shown in this table.

Q (cfs)	Maximum rate of INCREASE (cfs/day)	Maximum rate of DECREASE (cfs/day)
0-300	50	-50
301-650	75	-75
651-1400	150	-150
1401-3000	300	-600
>3000	1000	-2000

sfwmd.gov



## Other Considerations

- When possible, limit lake ascension rate in the Jun 1 - Aug 15 window to 0.25 ft per 7 days in Lakes Kissimmee, Cypress, Hatchineha (S-65), East Toho (S-59) and Toho (S-61).
- If outlook is for extreme dry conditions meet with KB staff to discuss modifications to this plan.

Slide Revised 7/29/2024

Figure KB-7. Headwaters Revitalization Schedule (HRS) Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A.

## **Lake Okeechobee**

Lake Okeechobee stage was 11.68 feet NAVD88 (12.98 ft NGVD29) on January 25, 2026, which was 0.10 feet lower than the previous week and 0.49 feet lower than a month ago (**Figure LO-1**). Lake stage is in Zone D of the regulation schedule, is 0.96 feet above the water shortage management band (**Figure LO-2**) and is 1.35 feet below the ecological envelope (**Figure LO-3**). According to NEXRAD, no rain fell directly over the lake during the previous week, and 0.80 inches were lost to evapotranspiration.

Average daily inflows (excluding rainfall) decreased from the previous week, dropping from 400 cfs to 320 cfs. The only notable inflows came from the Kissimmee River (320 cfs via S-65E(X1)). Average daily outflows (excluding evapotranspiration) increased slightly from the previous week, going from 2,020 cfs to 2,090 cfs. The highest release was to the west to the C-44 Canal through the S-77 structure (590 cfs). **Figures LO-4 and LO-5** show the combined average daily inflows and outflows for the lake over the past eight weeks, and average inflows and outflows last week, respectively.


In the most recent non-obsured satellite image from January 23, 2026, NOAA's Harmful Algal Bloom Monitoring System suggests the moderate to high cyanobacteria activity in the southern and western regions of the lake has returned after being temporarily dispersed by previous weeks strong winds and cooler temperatures (**Figure LO-6**).

The routine water quality and phytoplankton monitoring sampling trips are on the non-bloom season (Nov-Apr) once per month sampling schedule. Provisional phytoplankton results from the January 10-11 sampling showed 2 of 8 phytoplankton samples had detectable levels of microcystins, though both were  $< 0.5 \mu\text{g/L}$  and therefore well below EPA recreational standard (**Figure LO-7**). Five of the 30 water quality samples had chlorophyll *a*  $> 40 \mu\text{g/L}$ , 3 of which had values  $> 80 \mu\text{g/L}$  (all from the south end of the lake), indicating bloom level concentrations, while 12 samples had values between 20 and 40  $\mu\text{g/L}$  (**Figure LO-7**).

Note: All data presented in this report are provisional and are subject to change.


1 Month Ago:  
12/25/2025

Current:  
01/25/2026




**Figure LO-1.** Lake Okeechobee water depth estimates based on South Florida Water Depth Assessment Tool (SFWDAT).

## Lake Okeechobee Water Level History and Projected Stages




**Figure LO-2.** Recent Lake Okeechobee stages with projected stages based on a dynamic position analysis.  
Note: stages are in NGVD29, approximate NAVD88 values are shown for reference.


## Lake Okeechobee Stage vs Ecological Envelope



**Figure LO-3.** The current and seven prior year's annual lake stage hydrographs in comparison to the Lake Okeechobee ecological envelope (light grey).



**Figure LO-4.** Major inflows (orange) to and outflows east and west (blue) from Lake Okeechobee. Outflows south are shown in green. Flows into Lake Okeechobee from the L-8 canal through S-271 (formerly Culvert 10A) or from the C-44 canal through the S-308 are included as inflows. Conversely, flows from Lake Okeechobee into the L-8 or C-44 canals are included with outflows. Inflows are shown as positive values; outflows are negative. Outflows through the S-77 (Caloosahatchee) and S-308 (C-44 Canal) structures are based on downstream gauges to include flows to lock openings for navigation.



**Figure LO-5.** Inflows into Lake Okeechobee from Indian Prairie basins, Taylor Creek/Nubbin Slough, Kissimmee River and Fisheating Creek (currently no flow data available for FECR), and outflows to the west via S-77, to the east via S-308, to the south via S-351, S-352, S-354, and to southeast via S-271 (formerly Culvert 10A) for the week of Jan 19 - 25, 2026.

# Lake Okeechobee

## January 23, 2026


NOAA cyanobacteria product  
derived from Copernicus Sentinel-3  
OLCI data from EUMETSAT



True Color



Cyanobacteria  
Index




0 2.5 5  
Miles

**Figure LO-6.** Cyanobacteria bloom index level on Lake Okeechobee, based on NOAA's harmful algal bloom monitoring system. Gray color indicates cloud cover. \*Provisional NOAA image, subject to change\*.

Collection Date: January 10-11, 2026

Station	CHL $\alpha$ (ug/L)	TOXIN (ug/L)	TAXA	Station	CHL $\alpha$ (ug/L)	TOXIN (ug/L)	TAXA
FEBIN				L001	11.4		
FEBOUT				L004	10.2		
KISSR0.0	13.7	BDL	mixed	L006	14.3		
L005	17.4	BDL	Planktol	L007	44.3		
LZ2	23.1	BDL	mixed	L008	23.7		
KBARSE	7.4			LZ30	15.6	BDL	Microcys
RITTAE2	158.0	0.3	Micro/Raphi	LZ40	23.9		
PELBAY3	88.3			CLV10A	34.6	0.5	Microcys
POLE3S	40.6			NCENTER	13.2		
LZ25A	80.3			S308C	10.1		
PALMOUT	35.5	BDL	Micro/Raphi	S77	27.8		
PALMOUT1	29.8						
PALMOUT2	15.3						
PALMOUT3	16.5						
POLESOUT	14.8	BDL	Planktol				
POLESOUT1	21.0						
POLESOUT2	32.0						
POLESOUT3	25.2						
EASTSHORE	15.2						
NES135	21.2						
NES191	20.1						

Toxins include cylindrospermopsin and/or microcystins



**Figure LO-7.** Dominant taxa, cyanotoxins (ug/L) and chlorophyll  $\alpha$  (ug/L) concentration data from January 10 - 11, 2026. Sampling locations, chlorophyll  $\alpha$ , and total toxin concentrations are overlaid on the January 10, 2026, image from NOAA's harmful algal bloom monitoring system. Gray color indicates cloud cover.

## Estuaries

### ***St. Lucie Estuary***

Over the past week, mean total inflow to the St. Lucie Estuary was 110 cfs (**Figures ES-1** and **ES-2**), and the previous 30-day mean inflow was 124 cfs. For comparison, the historical provisional mean inflows from contributing areas are shown in **Figure ES-2**.

Over the past week, surface salinities increased slightly across all sites in the estuary (**Table ES-1** and **Figure ES-3**). The seven-day moving average of the surface and bottom salinities at the US1 Bridge was 26.5. Salinity conditions in the middle estuary were estimated to be within the upper stressed range for adult eastern oysters (**Figure ES-4**). The mean larval oyster recruitment rate reported by the Fish and Wildlife Research Institute (FWRI) for December was 0.3 spat/shell at Rio, which is a slight increase from the previous month (**Figure ES-5**).

### ***Caloosahatchee River Estuary***

Over the past week, mean total inflow to the Caloosahatchee River Estuary was 458 cfs (**Figures ES-6** and **ES-7**), and the previous 30-day mean inflow was 490 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-7**.

Over the past week, surface salinities increased at Shell Point and Sanibel and decreased at the remaining sites in the estuary (**Table ES-2** and **Figures ES-8** and **ES-9**). The seven-day mean salinities (**Table ES-2**) were in the optimal range (0-10) for tape grass at S-79 and Val I-75 and in the stressed range at Fort Myers. The seven-day mean salinity values were within the optimal range for adult eastern oysters at Cape Coral and in the upper stressed range at Shell Point and Sanibel (**Figure ES-10**). The mean larval oyster recruitment rates reported by the FWRI in December were 0 spat/shell at Iona Cove and 0.4 spat/shell at Bird Island, which is a decrease from the previous month (**Figures ES-11** and **ES-12**).

Surface salinity at Val I-75 was forecast for the next two weeks using an autoregression model (Qiu and Wan, 2013<sup>1</sup>) coupled with a linear reservoir model for the tidal basin. Model scenarios included pulse releases at S-79 ranging from 450 to 2,000 cfs, with estimated tidal basin inflows of 59 cfs. Model results from all scenarios predict daily salinity to be 7.0 or lower and the 30-day moving average surface salinity to be 8.2 or lower at Val I-75 at the end of the two-week period (**Table ES-3** and **Figure ES-13**). This keeps predicted salinities in the upper estuary within the optimal salinity range (0-10) for tape grass.

---

<sup>1</sup> Qui, C., and Y. Wan. 2013. Time series modeling and prediction of salinity in the Caloosahatchee River Estuary. *Water Resources Research* 49:5804-5816.

### **Red Tide**

The FWRI reported on January 23, 2026, that *Karenia brevis*, the Florida red tide dinoflagellate, was not observed in any samples collected within the District region over the past week.


### **Water Management Recommendations**

Lake stage is in Zone D. Current climatological and hydrological conditions are normal. The LOSOM release guidance suggests up to 2,000 cfs release at S-79 to the Caloosahatchee River Estuary and no releases at S-80 to the St. Lucie Estuary.


### **Minimum Flows and Minimum Levels**

The minimum flows and levels (MFL) for the Caloosahatchee River Estuary is a 30-day moving average flow of 457 cfs or greater at S-79. The current 30-day average flow at S-79 is 420 cfs (**Figure ES-14**) which is an exceedance.

The MFL for the Northwest Fork of the Loxahatchee River is a) flows at Lainhart Dam maintained at 35 cfs or greater and b) the 20-day moving average salinity of 2 or less at River Mile (RM) 9.2. An exceedance occurs when flows decline below 35 cfs for more than 20 consecutive days or when the 20-day moving average salinity at River Mile 9.2 exceeds 2. The current daily average flow at Lainhart Dam is 37 cfs (**Figure ES-15**) and the 20-day average salinity at RM 9.2 is 2.98, which is an exceedance (**Figure ES-16**).




**Figure ES-1.** Basins, water control structures and salinity monitoring sites in the St. Lucie Estuary.




**Figure ES-2.** Total daily inflows from Lake Okeechobee and runoff from the C-44, C-23, C-24, Ten Mile Creek, and Tidal Basins into the St. Lucie Estuary.

**Table ES-1.** Seven-day mean salinity at oyster monitoring sites in the St. Lucie Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope reflects the optimum salinity range for adult eastern oysters (*Crassostrea virginica*) in the estuary. Data are provisional.


Sampling Site	Surface	Bottom	Optimum Envelope
HR1 (North Fork)	<b>23.2</b> (22.6)	<b>25.4</b> (24.9)	10.0 – 25.0
US1 Bridge	<b>26.4</b> (26.0)	<b>26.6</b> (26.1)	10.0 – 25.0
A1A Bridge	<b>31.4</b> (31.0)	<b>31.7</b> (31.7)	10.0 – 25.0




**Figure ES-3.** Mean daily salinity at the A1A, US1, and HR1 sites in the St. Lucie Estuary.




**Figure ES-4.** Seven-day moving average of the surface and bottom salinities at the US1 Bridge in the St. Lucie Estuary.



**Figure ES-5.** Mean oyster recruitment at the Rio oyster monitoring station and 14-day mean salinity at US1 Bridge.




**Figure ES-6.** Basins, water control structures, and salinity monitoring sites in the Caloosahatchee River Estuary.




**Figure ES-7.** Total daily inflows from Lake Okeechobee, and runoff from the C-43 and Tidal basins into the Caloosahatchee River Estuary.

**Table ES-2.** Seven-day mean salinity at six monitoring sites in the Caloosahatchee River Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope in the upper estuary sites is for the protection of tape grass and the envelope in the lower estuary is the optimum salinity range for adult eastern oysters (*Crassostrea virginica*). Data are provisional.


Sampling Site	Surface	Bottom	Optimum Envelope
S-79 (Franklin Lock)	<b>5.9</b> (7.0)	<b>6.8</b> (7.6)	0.0 – 10.0
Val I-75	<b>7.4</b> (8.9)	<b>9.5</b> (9.8)	0.0 – 10.0
Fort Myers Yacht Basin	<b>14.8</b> (15.1)	<b>17.4</b> (16.1)	0.0 – 10.0
Cape Coral	<b>22.2</b> (22.3)	<b>23.3</b> (23.0)	10.0 – 25.0
Shell Point	<b>31.1</b> (30.6)	<b>31.5</b> (30.9)	10.0 – 25.0
Sanibel	<b>32.9</b> (32.4)	<b>33.0</b> (32.5)	10.0 – 25.0




**Figure ES-8.** Mean daily salinity at upper Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




**Figure ES-9.** Mean daily surface salinity at lower Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.



**Figure ES-10.** Seven-day moving average of surface and bottom salinities at Cape Coral, Shell Point, and Sanibel monitoring sites in the Caloosahatchee River Estuary.




**Figure ES-11.** Mean oyster recruitment at the Iona Cove oyster monitoring station and 14-day mean salinity at Cape Coral.




**Figure ES-12.** Mean oyster recruitment at the Bird Island oyster monitoring station and 14-day mean salinity at Shell Point.

**Table ES-3.** Predicted salinity at Val I-75 in the Caloosahatchee River Estuary at the end of the forecast period for various S-79 flow release scenarios.


Scenario	Simulated S-79 Flow (cfs)	Tidal Basin Runoff (cfs)	Daily Salinity	30-Day Mean Salinity
A	450	59	7.0	8.2
B	750	59	5.4	7.5
C	1,000	59	3.9	7.1
D	1,500	59	2.0	6.4
E	2,000	59	1.0	6.0




**Figure ES-13.** Surface salinity forecast at the Val I-75 site assuming a 450 cfs pulse release at S-79.



**Figure ES-14.** 30-day moving average flow at S-79 for the Caloosahatchee River Estuary Minimum Flows and Minimum Levels (MFL).



**Figure ES-15.** Average daily flow (cfs) at Lainhart Dam for the Loxahatchee River Estuary Minimum Flows and Minimum Levels (MFL).



**Figure ES-16.** 20-day moving average salinity at Lainhart Dam for the Loxahatchee River Estuary Minimum Flows and Minimum Levels (MFL).

## **Stormwater Treatment Areas**

**STA-1E:** STA-1E Central Flow-way is offline for construction activities. An operational restriction is in place in the Western Flow-way for post-construction vegetation grow-in. Online treatment cells are at or slightly above target stage. The 365-day PLR for the Western and Eastern Flow-way is below 1.0 g/m<sup>2</sup>/year (**Figure S-2**).

**STA-1W:** STA-1W Eastern Flow-way is offline for vegetation management activities. Most treatment cells are at target stage. Vegetation in the Western and Eastern Flow-ways is highly stressed. The 365-day PLRs for the Northern and Western Flow-ways are below 1.0 g/m<sup>2</sup>/year (**Figure S-2**).

**STA-2:** Operational restrictions are in place in Flow-ways 2, 3 and 4 for vegetation management activities. Treatment cells are at target stage or slightly below target stage. The 365-day PLRs for all Flow-ways are below 1.0 g/m<sup>2</sup>/year (**Figure S-3**).

**STA-3/4:** An operational restriction is in place in the Eastern Flow-way for vegetation management activities. Most treatment cells are slightly above target stage. Vegetation in the Central Flow-way is highly stressed. The 365-day PLR for the Eastern, Central, and Western Flow-ways are below 1.0 g/m<sup>2</sup>/year (**Figure S-3**).

**STA-5/6:** Treatment cells are at or above target stage. All treatment cells have highly stressed vegetation conditions. The 365-day PLRs for all Flow-ways are below 1.0 g/m<sup>2</sup>/year. (**Figure S-4**).

For definitions on STA operational language see glossary following figures


# Everglades Stormwater Treatment Areas - STAs

- Total WY2026 inflows to STAs (5/1/2025 to 1/25/2026): ~551,000 ac-ft
- Lake Okeechobee releases to FEBs/STAs
  - 1/19/2025 to 1/25/2026: 6,500 ac-ft
  - WY2026: ~ 48,100 ac-ft
- Extensive vegetation management activities underway to address stressed and highly stressed vegetation in EAV cells
- All treatment cells are at or near target water depth

Estimated Inflow and Outflow Volumes


Jan. 19<sup>th</sup>, 2026 - Jan 25<sup>th</sup>, 2026 *Includes preliminary data*

	Total Inflow (acre-feet)	Total Outflow (acre-feet)
STA-1E	0	60
STA-1W	3,300	70
STA-2	400	100
STA-3/4	1,500	1,000
STA-5/6	300	40



**Figure S-1.** STA depths and flow volumes

0 CFS Lake release capacity in Eastern Flow Path:  
1/26/2026 to 2/1/2026  
Subject to change weekly as dry season progresses





**Figure S-2.** Eastern Flow Path Weekly Status Report


700 CFS Lake release capacity in Central Flow Path:

1/26/2026 to 2/1/2026

- Subject to change weekly as dry season progresses



**Figure S-3.** Central Flow Path Weekly Status Report



**Figure S-4.** Western Flow Path Weekly Status Report

## Basic Concepts and Definitions for STA Weekly Status Report

- **Inflow:** Sum of flow volume at all inflow structures to an STA.
- **Lake Inflow:** Portion of the STA total inflow volume that originates from Lake Okeechobee.
- **Outflow:** Sum of flow volume at outflow structures from an STA.
- **Total Phosphorus (TP):** Total mass of phosphorus in all its forms; including particulate, dissolved, etc.
- **Inflow Concentration:** TP concentration is the mass of TP in micrograms per liter of water,  $\mu\text{g/L}$  or ppb. Inflow concentration refers to the flow-weighted mean TP from all inflow structures over a period of time.
- **Outflow Concentration:** The flow-weighted mean TP from all outflow structures over a period of time. The outflow concentration represents the reduction of inflow TP achieved by STA treatment of the inflow water.
- **WQBEL:** The STA outflow concentration that is required upon completion of the Restoration Strategies projects by December 2025. The outflow concentration shall not exceed 13 ppb as an annual flow weighted mean in more than 3 out of 5 water years on a rolling basis and shall not exceed 19 ppb as an annual flow weighted in any water year.
- **Flow-Way (FW):** One or more treatment cells connected in series. Cells typically have emergent aquatic vegetation (EAV) in the front portion of the flow-way followed by a mix of EAV and submerged aquatic vegetation (SAV).
- **Vegetation Status:** Healthy means the vegetation condition is good and will allow the STA to perform as designed. Stressed means the vegetation is showing signs of poor health, such as browning or areas of vegetation die-off, or the cell contains undesirable vegetation such as floating exotic vegetation requiring treatment. The TP reduction capability of the STA is affected when the vegetation condition is poor.
- **Phosphorus Loading Rate (PLR):** Mass of inflow TP in grams, divided by total treatment area of STA in square meters, per year. In general, a 365-day value of less than 1.0 is needed for an STA to perform optimally. A PLR of 2.0 is considered very high and a PLR of 3.0 is considered extremely high. The TP reduction capability of the STA is affected when the PLR is high, very high and extremely high.
- **Online:** Online status means the FW can receive and treat inflow.
- **Online with Restriction:** The FW can receive and treat inflow, but the amount of flow or water level may be limited temporarily. For example, a vegetation rehabilitation effort may require reduced flows through an area while the new plants are establishing, or nesting by protected species may require a certain water level not to be exceeded.
- **Offline:** The FW is unable to receive and treat inflow due to repairs, construction, or other prohibitive reasons.
- **Depth:** Difference between the average surface water level in a cell and the average ground elevation in that cell. Target depths, or depths between flow events, are between 1.25 ft to 1.5 ft. As depth approaches or drops below zero, an increasing percentage of the cell is considered dry and STA conditions deteriorate. An increase in depth above target depth is expected with increasing flow. However, as depth increases much above the target depth and is sustained over a period of time, it can be detrimental to vegetation health and overall STA treatment performance.
- **Note:** The data provided in this summary report were developed using a combination of provisional and quality-assured flow and water quality data. In some cases, best professional judgment was used to estimate missing data and revise questionable data. Values provided are not considered final but are appropriate for use in STA operational decision-making.

## **Everglades**

### ***Water Conservation Area Regulation Schedules***

WCA-1: Stage change at the 1-8C gauge last week remained at a steady recession, and stages were 0.55 feet below the falling A1 zone regulation line on Sunday, January 26<sup>th</sup>, 2026. WCA-2A: Last week's stage recession at the 2-17 gauge remains at a steady downward trend and is well above the falling zone A regulation line, at around 1.8 feet above on Sunday. WCA-3A: The 3-gauge average remains well in zone B and remains on a downward trend; on Sunday stages were 1.45 feet below the zone A regulation line. WCA-3A Northwest: Stage at Gauge 62 (NW corner) continued a steady decline last week, below the Upper Schedule regulation line by 0.89 feet on Sunday. See figures **EV-1** through **EV-4**.

### ***Water Depths***

The SFWDAT model output for January 25, 2026, illustrates recessions across the EPA. Over the past month there has been a slow recession in WCA-1, drying down to near soil surface in the north. The southern half of WCA-2A remains very deep for this time of year. WCA-2B is increasing in depth more so as increased pumping into this region from WCA-2A for water storage has begun. Drier than normal conditions expand across Northern WCA-3A. Depths are decreasing in WCA-3A and -3B and remain very low for this time of year with potential impacts on system-wide ecology. Hydrologic connectivity within the major sloughs of Everglades National Park has been declining; Shark River Slough has lost surface water connection going to the southwest coast while Taylor Slough may still have some hydrologic connectivity. Comparing current conditions to depths over the last twenty years; the majority of WCA-3A and WCA-3B are below the 10<sup>th</sup> percentiles (as they have for most of the last six months), while in southern WCA-2A and 2B depths remain above the 90th percentile. Depths throughout most of ENP have fallen to the 10<sup>th</sup> or 20<sup>th</sup> percentiles and WCA-1 depths have fallen into the 30<sup>th</sup> percentile or below. See figures **EV-5** through **EV-6**.

### ***Taylor Slough and Florida Bay***

All stages across Taylor Slough decreased over the past week, with an average decrease of 0.07 feet for the week. Changes ranged from -0.11 feet at E112 in the northern slough to -0.04 feet at P37 in the southern slough (**Figure EV-7** and **Figure EV-8**). Taylor Slough water levels remain below the recent average (WY1993-2016) for this time of year by 6.5 inches compared to before the Florida Bay Initiative (starting in 2017), an increase of 0.1 relative to January 15<sup>th</sup>. Stage at Taylor Slough Bridge (TSB) remains below ground, indicating a lack of water at the head of the slough (**Figure EV-8**). The Craighead Pond (CP) and TSB stages remain below the estimated average for 1900 by 0.75 and 2.42 feet, respectively.

Average Florida Bay salinity was 32.7, an increase of 3.9 from January 15<sup>th</sup>. Salinity changes ranged from +0.1 at Whipray Basin (WB) in the central offshore region to +8.5 at Terrapin Bay (TB) in the central nearshore region (**Figure EV-7**). Salinity is above the estimated average for 1900 and above the WY2001-2016 Interquartile Range (IQR) 75<sup>th</sup>

percentile in the eastern and central regions, and at the 75<sup>th</sup> percentile in the western region (**Figure EV-9**). Bay-wide salinity is above its recent average (WY1993-2016) for this time of year by 5.4, an increase of 2.1 from January 15<sup>th</sup>.

Salinity at the Taylor River (TR) station in the mangrove zone (tracked for the Florida Bay MFL) was 3.9, an increase of 2.7 from January 15<sup>th</sup> (**Figure EV-10**). The 365-day moving sum of flow from the five major creeks (McCormick Creek, Taylor River, Mud Creek, Trout Creek, West Highway Creek) was unable to be assessed due to missing data.

Average rainfall across Taylor Slough and Florida Bay was approximately 0.16 inches January 16<sup>th</sup> to 25<sup>th</sup>, based on the 18 gauges used for this report. Rainfall ranged from 0.02 inches at Garfield Bight (GB) in the western nearshore region to 0.38 inches at Manatee Bay (MB) in the eastern nearshore region (**Figure EV-11**). Wind directions and speeds in Florida Bay ranged from 0.4 mph SE on January 18<sup>th</sup> to 28.1 mph N on January 20<sup>th</sup> (**Figure EV-11**).

The Taylor River, Mud Creek, and West Highway Creek flow stations are currently offline until at least February, so data from all five major creeks are unable to be assessed. Based on the available data from Trout and McCormick Creeks, average daily flow totaled -2 acre-feet, with net negative flows for January 16<sup>th</sup> through 25<sup>th</sup>. Total daily creek flow ranged from -1,155 acre-feet on January 23<sup>rd</sup> to 1,258 acre-feet on January 19<sup>th</sup> (**Figure EV-13**). Average daily flow from Alligator Creek was 20 acre-feet, with net positive flows for the reporting period (**Figure EV-13**).


#### ***Implications/considerations for water management.***

- Stage recessions increased last week but would need to be reduced in order to protect the wetland ecology from damaging dry downs expected by the end of the dry season in most regions.
  - The depths within WCA-3A this wet season were not high enough to recover aquatic prey populations from antecedent dry conditions or protect peat soils throughout the current dry season.
  - WCA-3A continues to experience unseasonably dry conditions.
  - This has the potential to further extend the recent run of 4 consecutive poor wading bird nesting years into the 2026 nesting seasons.
  - With La Nina conditions this dry season, conserving water within the WCAs will continue to be ecologically beneficial, especially in regions prone to dry out (e.g. WCA-3A North).
- Depths are too deep (<2.15 feet) in south-central WCA-2A where shallower conditions (read suitable depth) are needed to recover ridge and slough habitat.
- Taylor Slough depths are receding quicker than usual for this time of year, slough connectivity is almost lost, and salinities are above their recent average in Florida Bay.


- Freshwater input into Taylor Slough and the C-111 basin would help moderate salinities and support recovery of estuarine conditions in Florida Bay.
- With flows across Shark River Slough now more restricted, conserving water within WCA-3A and prioritizing southern deliveries through Taylor Slough may provide greater ecological benefits at the broader ecosystem scale. Conserving water in the WCAs while providing freshwater input to the sloughs of ENP will require careful consideration of a balance between the upstream and downstream ecological needs of the system. Individual regional recommendations can be found in **Table EV-2**.

**Table EV-2.** Previous week's rainfall and water depth changes in Everglades basins.


Everglades Region	Rainfall (inches)	Stage change (feet)
WCA-1	0.15	-0.09
WCA-2A	0.07	-0.16
WCA-2B	0.05	-0.02
WCA-3A	0.04	-0.11
WCA-3B	0.03	-0.07
ENP	0.15	-0.06




**Figure EV-1.** WCA-1 stage hydrographs and regulation schedule.




**Figure EV-2.** WCA-2A stage hydrographs and regulation schedule.




**Figure EV-3.** WCA-3A stage hydrographs (three-gauge average, 3-69W) and regulation schedule.




**Figure EV-4.** WCA-3A stage hydrograph (Deer gauge; Site 62) and regulation schedule.




**Figure EV-5.** Everglades water depths from two months ago (left), one month ago (center) and present (right), based on SFWDAT.




**Figure EV-6.** Present water depths (January 25, 2026) compared to the day of year relative to average (percentile) over the previous 20 years.




**Figure EV-7.** Taylor Slough water depths and Florida Bay salinities with changes since January 15<sup>th</sup>.




**Figure EV-8.** Taylor Slough water depth time series for Taylor Slough Bridge (TSB; northern slough) and Craighead Pond (CP; southern slough).




**Figure EV-9.** Eastern (top panel), Central (middle panel) and Western (bottom panel) Florida Bay daily average salinities with WY2001-2016 interquartile (25-75 percentile) ranges (IQR) and estimated historical daily average salinities. The hypersalinity threshold indicates the level at which salinities start to become harmful to seagrass.



**Figure EV-10.** Daily average salinity at Taylor River (TR) tracked for the Florida Bay MFL criteria. The 365-day total creek flow MFL metric is not currently available due to missing creek flow data.



**Figure EV-11.** Daily average rain across Taylor Slough and Florida Bay, along with hourly average wind speed and direction (measured at Long Key) in Florida Bay over the past four weeks.



**Figure EV-12.** Top: daily average creek flow summed between the five major creeks with estimated historical daily flow over the past four weeks (**note**: data from Taylor River, Mud Creek and West Highway Creek are currently unavailable since November 21<sup>st</sup>, 2025). Bottom: Daily average Alligator Creek flow data. N/A indicates missing data.

**Table EV-2.** Weekly water depth changes and water management recommendations

<b>SFWMD Everglades Ecological Recommendations, January 26, 2026 (red is new)</b>			
	Weekly change	Recommendation	Reasons
<b>WCA-1</b>	<b>Stage decreased by 0.09 feet.</b>	A recession of no faster than 0.05 feet per week.	Conserve water, maintain within basin and downstream habitat and wildlife. Maintain maintenance access for vegetation management.
<b>WCA-2A</b>	<b>Stage decreased by 0.16 feet</b>	A recession of no faster than 0.12 feet per week.	Maintain within basin (north versus south) and downstream habitat and wildlife.
<b>WCA-2B</b>	<b>Stage increased by 0.02 feet</b>	A recession of no faster than 0.12 feet per week.	Protect within basin and downstream habitat and wildlife.
<b>WCA-3A NE</b>	<b>Stage decreased by 0.14 feet</b>	A recession of no faster than 0.05 feet per week.	Conserve water, maintain within basin and downstream habitat and wildlife. Provide suitable depths for aquatic prey and protect against peat soil loss during the dry season.
<b>WCA-3A NW</b>	<b>Stage decreased by 0.08 feet</b>	A recession of no faster than 0.05 feet per week.	Conserve water, maintain within basin and downstream habitat and wildlife. Provide suitable depths for aquatic prey and protect against peat soil loss during the dry season.
<b>Central WCA-3A S</b>	<b>Stage decreased by 0.12 feet</b>	A recession of no faster than 0.05 feet per week.	Conserve water, maintain within basin and downstream habitat and wildlife. Provide suitable depths for aquatic prey and protect against peat soil loss during the dry season.
<b>Southern WCA-3A S</b>	<b>Stage decreased by 0.10 feet</b>		
<b>WCA-3B</b>	<b>Stage decreased by 0.07 feet</b>	A recession of no faster than 0.12 feet per week.	Protect within basin and downstream habitat and wildlife.
<b>ENP-SRS</b>	<b>Stage decreased by 0.06 feet.</b>	Make discharges to ENP according to COP protocol, considering up/down stream ecological conditions.	Protect within basin and upstream habitat and wildlife.
<b>Taylor Slough</b>	<b>Stage changes ranged from -0.11 feet to -0.04 feet</b>	Move water southward as possible.	When available, provide freshwater to promote water movement.
<b>FB- Salinity</b>	<b>Salinity changes ranged from +0.1 to +8.5</b>	Move water southward as possible.	When available, provide freshwater to promote water movement.