

Disclaimer: Information contained in the report addresses environmental conditions only and is not the official South Florida Water Management District operations recommendation or decision.

M E M O R A N D U M

TO: John Mitnik, Assistant Executive Director, Executive Office Staff

FROM: SFWMD Staff Environmental Advisory Team

DATE: April 15, 2026

SUBJECT: Weekly Environmental Conditions for Systems Operations

Summary

Weather Conditions and Forecast

Wednesday a few light showers along and near the east coast and through the Florida Keys are possible, while temperatures gradually rise under a breezy easterly to east-northeasterly flow, particularly south and east of Lake Okeechobee. The combination of dry conditions, persistent breeziness, warming temperatures, and the strong April sun angle will promote elevated evapotranspiration rates across the region. Increased moisture could support scattered shower activity moving onshore into the east-coast basins, the Florida Keys, and portions of the eastern interior beginning Friday morning and continuing into the afternoon, with additional activity possible overnight. Even so, area-averaged rainfall is expected to remain limited, with only isolated locations — primarily near the coast — potentially receiving locally higher amounts. The moisture field will expand spreading farther inland and supporting additional shower development from Lake Okeechobee southward—particularly during the morning and afternoon hours on Saturday—although overall rainfall should remain modest and largely confined to the southeastern third to half of the SFWMD. Temperatures will also warm more quickly, and western areas are likely to see near-record or record heat. A cold front will move into the northern part of the SFWMD overnight Sunday and the southern part of the SFWMD by Monday morning. The cold front should gradually slide southward through the remainder of the SFWMD by Monday evening. Elevated moisture levels along and ahead of the front, combined with an increase in atmospheric instability, are expected to support increased rainfall on Monday, with the highest area-averaged rainfall most likely along and near the east coast. Given the anticipated slowing of the front, deeper moisture could persist into Tuesday, which would support additional rainfall, especially across the east-coast basins in what could be a mini east-coast rain event. Behind the cold front, a marked cooling trend is expected, accompanied by very breezy northeasterly winds that will gradually shift east-northeasterly and easterly by Tuesday, followed by drier air expanding across most of the SFWMD. For the week ending next Tuesday morning, total area-averaged rainfall across the SFWMD is expected to be below to well below the long-term average, despite the increase in rainfall forecast late in the period.

Kissimmee

In the past week, releases were made as needed from East Lake Toho and Lake Toho to continue snail kite nesting season stage recessions to reach low pool by June 1, 2026. Releases from Lakes Kissimmee-Cypress-Hatchineha followed the Headwaters Revitalization Schedule Increment 1 Temporary Deviation Discharge Plan. Weekly average discharge on April 12, 2026, was 590 cfs at S-65 and 520 cfs at S-65A. Mean weekly water depth on the Kissimmee River floodplain was unchanged from the previous week's value of 0.33 feet. The weekly average concentration of dissolved oxygen in the Kissimmee River decreased from 8.1 mg/L the previous week to 7.4 mg/L, which is above both the potentially lethal level of 1.0 mg/L and the stressful level of 2.0 mg/L for Florida bass and other species.

Lake Okeechobee

Lake Okeechobee stage was 10.77 feet NAVD88 (12.08 ft NGVD29) on April 12, 2026, which was 0.07 feet higher than the previous week and 0.07 feet lower than a month ago. Average daily inflows (excluding rainfall) decreased from 750 cfs the previous week to 630 cfs. Average daily outflows (excluding evapotranspiration) rose from 10 cfs to 20 cfs. The most recent non-obscured satellite image from April 12, 2026, NOAA's Harmful Algal Bloom Monitoring System suggests moderate to high cyanobacteria potential in Fisheating Bay.

Estuaries

Total inflow to the St. Lucie Estuary averaged 2,837 cfs over the past week with no flow coming from Lake Okeechobee. Mean salinities decreased at all sites over the past week. Salinity in the middle estuary was in the optimal range (10-25) for adult eastern oysters.

Total inflow to the Caloosahatchee River Estuary averaged 2,009 cfs over the past week with no flow coming from Lake Okeechobee. Over the past week, salinities decreased at all sites in the estuary. Salinities were in the optimal range (0-10) for tape grass in the upper estuary. Salinities were within the optimal range for adult eastern oysters at Cape Coral and in the upper stressed range at Shell Point and Sanibel.

Stormwater Treatment Areas

For the week ending April 12, 2026, no Lake Okeechobee water was delivered to the FEBs/STAs. The total amount of Lake releases sent to the FEBs/STAs in WY2026 is approximately 97,400 ac-feet. The total amount of inflows to the STAs in WY2026 is approximately 677,000 ac-feet. Most online STA treatment cells are at or above target stage. STA-1E Central Flow-way is offline for construction activities. STA-1W Eastern Flow-way is offline for vegetation management activities. Operational restrictions are in effect in STA-1E Western Flow-way and STA-3/4 Eastern Flow-way for vegetation management activities. This week, if LOSOM recommends Lake releases to the WCAs and conditions allow, releases will be sent to STA-2.

Everglades

Above-average rainfall occurred across most of the Everglades Protection Area (EPA), with the highest totals in WCA-1 (where average stage rose 0.19 feet) and the lowest in WCA-3A (where stage in the central region has fallen by nearly 0.4 feet over the last two weeks). The trend in rainfall was reflected by the decrease in recession rates throughout the EPA; average stage change was positive with an ascension rate of 0.05 feet/week. Rainfall over the past few weeks has increased water depths across much of the EPA relative to the past 20 years, with the exception of WCA-3A and WCA-3B, which continue to exhibit well below-average depths. These conditions can have ecological consequences, including reduced already-limited prey populations, increased risk of damaging wildfires, enhanced peat oxidation, and potential ridge and slough degradation. Wading bird activity remains very limited with nesting in numbers only occurring in WCA-1, some foraging is occurring along the drying front in WCA-2A, and there is very little foraging or nesting occurring in WCA-3A. Wood storks have abandoned nests system wide. Unless we get an unexpected large increase in nesting in the next few weeks, this nesting season could be one of the lowest nesting efforts observed in the Everglades in the last 30 years. Taylor Slough stages decreased last week but remain above the recent averages for this time of year by 8 inches. Average Florida Bay salinity decreased slightly last week, and the central region is now just below the interquartile range.

Supporting Information

Kissimmee Basin

Upper Kissimmee

On April 12, 2026, mean daily lake stages were 54.9 feet NAVD88 (1.1 feet below schedule) in East Lake Toho, 52.0 feet NAVD88 (1.0 feet below schedule) in Lake Toho, and 48.8 feet NAVD88 (2.0 feet below the Increment 1 Temporary Deviation schedule) in Lakes Kissimmee-Cypress-Hatchineha (KCH) (**Table KB-1, Figures KB-1-3**).

Lower Kissimmee

For the week ending April 12, 2026, mean weekly discharge was 590 cfs at S-65 and 520 cfs at S-65A. Mean weekly discharge from the Kissimmee River was 490 cfs at S-65D and 430 cfs at S-65E (**Table KB-2**). Mean weekly headwater stages were 45.2 feet NAVD88 at S-65A and 29.1 feet NAVD88 at S-65D. Mean weekly river channel stage increased from 31.4 feet the previous week to 31.7 feet NAVD88 (**Figure KB-4**). Mean weekly water depth on the Kissimmee River floodplain was 0.33 feet, unchanged from the previous week (**Table KB-2, Figure KB-5**). The weekly average concentration of dissolved oxygen in the Kissimmee River decreased from 8.1 mg/L the previous week to 7.4 mg/L (**Table KB-2, Figure KB-6**).

Water Management Recommendations

Continue the stage recessions in East Lake Toho and Lake Toho to reach their low pools on June 1, 2026. In KCH, follow the Headwaters Revitalization Schedule (HRS) Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A (**Figure KB-7**). With KCH stage in Zone B3, target flows between 300 and 1,400 cfs at S-65A, using the Increment 1 Interpolation Tool to determine discharge relative to stage in KCH; if stage decreases into Zone B4, target flows of 300 cfs.

Table KB-1. Average discharge for the preceding seven days, Sunday’s average daily stage and Sunday’s average daily departure from Kissimmee Chain of Lakes (KCOL) flood regulation lines or temporary schedules. All data are provisional.

Water Body	Structure	Stage Monitoring Site	Weekly (7-Day) Average Discharge (cfs)	Sunday Lake Stage (feet NAVD88) ^a	Schedule Type ^b	Sunday Schedule Stage (feet NAVD88)	Sunday Departure from Regulation (feet)	
							4/12/26	4/5/26
Lakes Hart and Mary Jane	S-62	LKMJ	40	59.1	R	59.4	-0.3	-0.2
Lakes Myrtle, Preston and Joel	S-57	S-57	7	59.4	R	59.5	-0.1	0.0
Alligator Chain	S-60	ALLI	120	62.2	R	62.2	0.0	0.0
Lake Gentry	S-63	LKGT	180	59.7	R	59.7	0.0	0.0
East Lake Toho	S-59	TOHOE	290	54.9	R	56.0	-1.1	-1.1
Lake Toho	S-61	TOHOW S-61	740	52.0	R	53.0	-1.0	-1.1
Lakes Kissimmee, Cypress and Hatchineha	S-65	KUB011 LKIS5B	590	48.8	T	50.8	-2.0	-2.3

a. Names of in-lake monitoring sites and structures used to determine lake stage. If more than one site is listed, an average is reported.

b. A: projected recession line; R: USACE regulation schedule; S: temporary recession target line; T: temporary schedule; NA: not applicable or not available.




Figure KB-1. East Lake Toho regulation schedule, stage, discharge, and rainfall.




Figure KB-2. Lake Toho regulation schedule, stage, discharge, and rainfall.




Figure KB-3. Lakes Kissimmee, Cypress and Hatchineha regulation schedule, stage, discharge, and rainfall.

Table KB-2. One- and seven-day average discharge and stage at Lower Kissimmee basin structures, river channel dissolved oxygen concentrations and water depths in the Phase I area floodplain. All data are provisional.

Metric	Location	Sunday Daily Average	Weekly Average for Previous Seven Day Periods			
		4/12/26	4/12/26	4/5/26	3/29/26	3/22/26
Discharge	S-65	870	590	380	420	510
Discharge	S-65A ^a	760	520	340	360	450
Headwater Stage (feet NAVD88)	S-65A	45.2	45.2	45.3	45.4	45.2
Discharge	S-65D ^b	700	490	420	410	560
Headwater Stage (feet NAVD88)	S-65D ^c	24.6	29.1	28.8	28.8	29.4
Discharge (cfs)	S-65E ^d	570	430	340	320	420
Discharge (cfs)	S-67	0	0	0	0	0
Dissolved Oxygen (mg/L) ^e	Phase I, II/III river channel	8.4	7.4	8.1	8.6	8.5
River channel mean stage (feet NAVD88) ^f	Phase I river channel	32.6	31.7	31.4	31.3	32.0
Mean depth (feet) ^g	Phase I & II/III floodplain	0.34	0.33	0.33	0.33	0.33

a. Combined discharge from main and auxiliary structures.

b. Combined discharge from S-65D, S-65DX1, and S-65DX2.

c. Average stage from S-65D and S-65DX1.


d. Combined discharge from S-65E and S-65EX1.

e. Dissolved oxygen is the average of values from sondes KRBN, PC62, PC33, PD62R, and PD42R.

f. Mean of five river channel stations (PC62, KRDR02, KRBN, PC33, PC11) in the Phase I area.

g. One-day spatial average obtained from the South Florida Water Depth Assessment Tool (SFWDAT).

Stage and Discharge Hydrograph (report date: 4/14/2026; data through: 4/12/2026)



*River Channel Stage is the average for PC62, KRDR02, KRBN, PC33, and PC11.



Figure KB-4. Kissimmee River stage, discharge, and rainfall.





Figure KB-5. Phase I-II-III area Kissimmee River floodplain water depths (from left to right) one year ago, one month ago, and current.



Report Date: 4/14/2026; data are through: 4/12/2026



Figure KB-6. Kissimmee River channel mean daily dissolved oxygen concentration (mg/L), S-65A discharge (cfs), temperature (°C) and rainfall (inches). Dissolved oxygen (DO) and temperature are mean daily values averaged for PC62, KRDR02, KRBN, PC33, PC11, PD62R, and PD42R with an average of five stations reporting this week. Rainfall values are daily totals for Kissimmee River (Pool BCD) AHED watershed.

HRS Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A

Discharge Guidance for Increment 1 Temporary Deviation Discharge Plan		
ZONE	S-65 RELEASES	S-65A TARGET FLOWS
ZONE A	Releases for Flood Risk Management up to maximum structure capacity as determined by downstream constraints with a firm capacity of 3,000 cfs.	
ZONE B1	1,400 cfs minimum ramp to 3,000 cfs at Zone A boundary	S-65A releases between 1,400 cfs and 3,000 cfs at Zone A boundary based on Table 1
ZONE B2	Releases as needed to target flows at S-65A	Target S-65A flows of 1,400 cfs to meet ecological needs
ZONE B3	Releases as needed to target flows at S-65A	S-65A flows between 300 cfs and 1,400 cfs
ZONE B4	Releases as needed to target flows at S-65A	Target S-65A flows of 300 cfs
ZONE B5	Releases as needed to target flows at S-65A	Target S-65A flows of 150 cfs
ZONE C	0 cfs	Flow as needed to maintain optimum S-65A headwater




Table KB-3. Maximum Rate of Change Limits for S-65A

MAXIMUM Release Rate of Change Limits for S-65A. In general recommended rates of change will be slower than shown in this table.

Q (cfs)	Maximum rate of INCREASE (cfs/day)	Maximum rate of DECREASE (cfs/day)
0-300	50	-50
301-650	75	-75
651-1400	150	-150
1401-3000	300	-600
>3000	1000	-2000

CENTRAL AND SOUTHERN FLORIDA PROJECT
 2024 LAKE KISSIMMEE, HATCHINEHA & CYPRESS
 Temporary Deviation (Increment 1)
 DATED: May 2024
 DEPARTMENT OF THE ARMY, JACKSONVILLE DISTRICT
 CORPS OF ENGINEERS, JACKSONVILLE, FLORIDA

Other Considerations

- When possible, limit lake ascension rate in the Jun 1 - Aug 15 window to 0.25 ft per 7 days in Lakes Kissimmee, Cypress, Hatchineha (S-65), East Toho (S-59) and Toho (S-61).
- If outlook is for extreme dry conditions meet with KB staff to discuss modifications to this plan.

Figure KB-7. Headwaters Revitalization Schedule (HRS) Increment 1 Temporary Deviation Discharge Plan for S-65/S-65A.

Lake Okeechobee

Lake Okeechobee stage was 10.77 feet NAVD88 (12.08 ft NGVD29) on April 12, 2026, which was 0.07 feet higher than the previous week and 0.07 feet lower than a month ago (**Figure LO-1**). Lake stage is in Zone D of the regulation schedule, is 0.68 feet above the water shortage management band (**Figure LO-2**) and 0.52 feet below the ecological envelope (**Figure LO-3**). According to NEXRAD, 2.14 inches of rain fell directly over the lake during the previous week, and 1.15 inches were lost to evapotranspiration.

Average daily inflows (excluding rainfall) decreased from 750 cfs the previous week to 630 cfs. The highest average inflow came from the Kissimmee River (420 cfs via S-65E(X1)). Average daily outflows (excluding evapotranspiration) rose from 10 cfs to 20 cfs. The only release was to the west through the S-77 structure (20 cfs). **Figures LO-4 and LO-5** show the combined average daily inflows and outflows for the lake over the past eight weeks, and the average inflows and outflows last week, respectively.

In the most recent non-obscured satellite image from April 12, 2026, NOAA's Harmful Algal Bloom Monitoring System suggests moderate to high cyanobacteria potential in Fisheating Bay. (**Figure LO-6**).

Note: All data presented in this report are provisional and are subject to change.

1 Month Ago:
03/12/2026

Current:
04/12/2026

10.85 ft
NAVD88

10.77 ft
NAVD88




Figure LO-1. Lake Okeechobee water depth estimates based on South Florida Water Depth Assessment Tool (SFWDAT).




Figure LO-2. Recent Lake Okeechobee stages with projected stages based on a conditional position analysis. Note: stages are in NGVD29, approximate NAVD88 values are shown for reference.

Lake Okeechobee Stage vs Ecological Envelope




Figure LO-3. The current and select prior year's annual lake stage hydrographs in comparison to the Lake Okeechobee ecological envelope (light grey).




Figure LO-4. Major inflows (orange) to and outflows east and west (blue) from Lake Okeechobee. Outflows south are shown in green. Flows into Lake Okeechobee from the L-8 canal through S-271 (formerly Culvert 10A) or from the C-44 canal through the S-308 are included as inflows. Conversely, flows from Lake Okeechobee into the L-8 or C-44 canals are included with outflows. Inflows are shown as positive values; outflows are negative. Outflows through the S-77 (Caloosahatchee) and S-308 (C-44 Canal) structures are based on downstream gauges to include flows to lock openings for navigation.




Figure LO-5. Inflows into Lake Okeechobee from Indian Prairie basins, Taylor Creek/Nubbin Slough, Kissimmee River and Fisheating Creek (currently no flow data available for FECR), and outflows to the west via S-77, to the east via S-308, to the south via S-351, S-352, S-354, and to southeast via S-271 (formerly Culvert 10A) for the week of Apr 6 - 12, 2026.




Figure LO-6. Cyanobacteria bloom index level on Lake Okeechobee, based on NOAA’s harmful algal bloom monitoring system. Gray color indicates cloud cover. *Provisional NOAA image, subject to change*.

Estuaries

St. Lucie Estuary

Over the past week, mean total inflow to the St. Lucie Estuary was 2,837 cfs (**Figures ES-1 and ES-2**), and the previous 30-day mean inflow was 808 cfs. For comparison, the historical provisional mean inflows from contributing areas are shown in **Figure ES-2**.

Over the past week, salinities decreased at the A1A Bridge, HR1, and US1 Bridge sites (**Table ES-1 and Figure ES-3**). The seven-day moving average of the surface and bottom salinities at the US1 Bridge was 18.1. Salinity conditions in the middle estuary were estimated to be within the optimal range for adult eastern oysters (**Figure ES-4**). The mean larval oyster recruitment rate reported by the Fish and Wildlife Research Institute (FWRI) for March was 0 spat/shell at Rio, which is equal to the previous month (**Figure ES-5**). Oyster density at Rio was 464 oysters/m², which is slightly lower than the previous survey in December.

Caloosahatchee River Estuary

Over the past week, mean total inflow to the Caloosahatchee River Estuary was 2,009 cfs (**Figures ES-6 and ES-7**), and the previous 30-day mean inflow was 535 cfs. For comparison, the historical provisional mean inflows from the contributing areas are shown in **Figure ES-7**.

Over the past week, salinities decreased at all sites in the estuary (**Table ES-2 and Figures ES-8 and ES-9**). The seven-day mean salinities (**Table ES-2**) were in the optimal range (0-10) for tape grass in the upper estuary at S-79 and Val I-75 and in the damaging range (>15) at Fort Myers. The seven-day mean salinity values were within the optimal range for adult eastern oysters at Cape Coral and in the upper stressed range at Shell Point and Sanibel (**Figure ES-10**). The mean larval oyster recruitment rates reported by the FWRI in March were 0.03 spat/shell at Iona Cove and 2.9 spat/shell at Bird Island, which is an increase at both sites from the previous month (**Figures ES-11 and ES-12**). Oyster density was 784 oysters/m² at Iona Cove and 1,310 oysters/m² at Bird Island which are both slightly lower than the previous surveys in December.

Surface salinity at Val I-75 was forecast for the next two weeks using an autoregression model (Qiu and Wan, 2013¹) coupled with a linear reservoir model for the tidal basin. Model scenarios included pulse releases at S-79 ranging from 450 to 2,000 cfs, with estimated tidal basin inflows of 65 cfs. Model results from all scenarios predict daily salinity to be 5.3 or lower and the 30-day moving average surface salinity to be 6.3 or lower at Val I-75 at the end of the two-week period (**Table ES-3 and Figure ES-13**). This keeps predicted salinities in the upper estuary within the optimal salinity range (0-10) for tape grass.

¹ Qiu, C., and Y. Wan. 2013. Time series modeling and prediction of salinity in the Caloosahatchee River Estuary. *Water Resources Research* 49:5804-5816.

Red Tide

The FWRI reported on April 10, 2026, that *Karenia brevis*, the Florida red tide dinoflagellate, was not observed in any samples collected within the District region.

Water Management Recommendations

Lake stage is in Zone D. Current climatological and hydrological conditions are normal. The LOSOM release guidance suggests up to 2,000 cfs release at S-79 to the Caloosahatchee River Estuary and no releases at S-80 to the St. Lucie Estuary.

Minimum Flows and Minimum Levels

The minimum flows and levels (MFL) for the Caloosahatchee River Estuary is a 30-day moving average flow of 457 cfs or greater at S-79. The current 30-day average flow at S-79 is 875.9 cfs (**Figure ES-14**).

The MFL for the Northwest Fork of the Loxahatchee River is a) flows at Lainhart Dam maintained at 35 cfs or greater and b) the 20-day moving average salinity of 2 or less at River Mile (RM) 9.2. An exceedance occurs when flows decline below 35 cfs for more than 20 consecutive days or when the 20-day moving average salinity at River Mile 9.2 exceeds 2. The current daily average flow at Lainhart Dam is 105 cfs (**Figure ES-15**) and the 20-day average salinity at RM 9.2 is 0.7 (**Figure ES-16**).




Figure ES-1. Basins, water control structures and salinity monitoring sites in the St. Lucie Estuary.




Figure ES-2. Total daily inflows from Lake Okeechobee and runoff from the C-44, C-23, C-24, Ten Mile Creek, and Tidal Basins into the St. Lucie Estuary.

Table ES-1. Seven-day mean salinity at oyster monitoring sites in the St. Lucie Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope reflects the optimum salinity range for adult eastern oysters (*Crassostrea virginica*) in the estuary. Data are provisional.

Sampling Site	Surface	Bottom	Optimum Envelope
HR1 (North Fork)	11.0 (21.4)	20.6 (22.3)	10.0 – 25.0
US1 Bridge	17.1 (25.3)	19.1 (25.7)	10.0 – 25.0
A1A Bridge	25.9 (31.4)	28.3 (31.7)	10.0 – 25.0




Figure ES-3. Mean daily salinity at the A1A, US1, and HR1 sites in the St. Lucie Estuary.




Figure ES-4. Seven-day moving average of the surface and bottom salinities at the US1 Bridge in the St. Lucie Estuary.




Figure ES-5. Mean oyster recruitment at the Rio oyster monitoring station and 14-day mean salinity at US1 Bridge.




Figure ES-6. Basins, water control structures, and salinity monitoring sites in the Caloosahatchee River Estuary.




Figure ES-7. Total daily inflows from Lake Okeechobee, and runoff from the C-43 and Tidal basins into the Caloosahatchee River Estuary.

Table ES-2. Seven-day mean salinity at six monitoring sites in the Caloosahatchee River Estuary. Current means are in bold font; previous week's means are in parentheses. The envelope in the upper estuary sites is for the protection of tape grass and the envelope in the lower estuary is the optimum salinity range for adult eastern oysters (*Crassostrea virginica*). Data are provisional.

Sampling Site	Surface	Bottom	Optimum Envelope
S-79 (Franklin Lock)	3.6 (7.0)	3.7 (7.2)	0.0 – 10.0
Val I-75	4.8 (8.3)	6.4 (10.2)	0.0 – 10.0
Fort Myers Yacht Basin	12.7 (16.5)	17.5 (18.4)	0.0 – 10.0
Cape Coral	20.4 (22.6)	21.8 (24.3)	10.0 – 25.0
Shell Point	30.2 (32.7)	31.2 (33.7)	10.0 – 25.0
Sanibel	34.1 (35.0)	34.9 (36.7)	10.0 – 25.0




Figure ES-8. Mean daily salinity at upper Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




Figure ES-9. Mean daily surface salinity at lower Caloosahatchee River Estuary monitoring sites and mean daily flow at S-79.




Figure ES-10. Seven-day moving average of surface and bottom salinities at Cape Coral, Shell Point, and Sanibel monitoring sites in the Caloosahatchee River Estuary.




Figure ES-11. Mean oyster recruitment at the Iona Cove oyster monitoring station and 14-day mean salinity at Cape Coral.




Figure ES-12. Mean oyster recruitment at the Bird Island oyster monitoring station and 14-day mean salinity at Shell Point.

Table ES-3. Predicted salinity at Val I-75 in the Caloosahatchee River Estuary at the end of the forecast period for various S-79 flow release scenarios.

Scenario	Simulated S-79 Flow (cfs)	Tidal Basin Runoff (cfs)	Daily Salinity	30-Day Mean Salinity
A	450	65	5.3	6.3
B	750	65	4.1	5.8
C	1,000	65	3.0	5.4
D	1,500	65	1.6	5.0
E	2,000	65	0.7	4.6

Observed and Forecasted Flow at S-79 and Salinity at Val I-75




Figure ES-13. Surface salinity forecast at the Val I-75 site assuming a 450 cfs pulse release at S-79.




Figure ES-14. 30-day moving average flow at S-79 for the Caloosahatchee River Estuary Minimum Flows and Minimum Levels (MFL).




Figure ES-15. Average daily flow (cfs) at Lainhart Dam for the Loxahatchee River Estuary Minimum Flows and Minimum Levels (MFL).




Figure ES-16. 20-day moving average salinity at Lainhart Dam for the Loxahatchee River Estuary Minimum Flows and Minimum Levels (MFL).

Stormwater Treatment Areas

STA-1E: STA-1E Central Flow-way is offline for construction activities. An operational restriction is in place in the Western Flow-way for post-construction vegetation grow-in. Online treatment cells are slightly above target stage. The 365-day PLR for the Western and Eastern Flow-way is below 1.0 g/m²/year (**Figure S-2**).

STA-1W: STA-1W Eastern Flow-way is offline for vegetation management activities. Treatment cells are at or slightly above target stage. Vegetation in the Western and Eastern Flow-ways is highly stressed. The 365-day PLRs for the Northern and Western Flow-ways are below 1.0 g/m²/year (**Figure S-2**).

STA-2: Treatment cells are at or slightly above target stage. The 365-day PLRs for all Flow-ways are below 1.0 g/m²/year (**Figure S-3**).

STA-3/4: An operational restriction is in place in the Eastern Flow-way for vegetation management activities. Most treatment cells are at or slightly above target stage. Vegetation in the Central Flow-way is highly stressed. The 365-day PLR for the Eastern, Central, and Western Flow-ways are below 1.0 g/m²/year (**Figure S-3**).

STA-5/6: Treatment cells are at or below target stage. All treatment cells have highly stressed vegetation conditions. The 365-day PLRs for all Flow-ways are below 1.0 g/m²/year. (**Figure S-4**).

For definitions on STA operational language see glossary following figures

Everglades Stormwater Treatment Areas - STAs

- Total WY2026 inflows to STAs (5/1/2025 to 4/12/2026): ~677,000 ac-ft
- Lake Okeechobee releases to FEBs/STAs
 - 4/6/2026 to 4/12/2026: 0 ac-ft
 - WY 2026: ~ 97,400 ac-ft
- Extensive vegetation management activities underway to address stressed and highly stressed vegetation in EAV cells
- All treatment cells are at or near target water depth

Estimated Inflow and Outflow Volumes

Apr. 6th, 2026 - Apr. 12th, 2026 *Includes preliminary data*

	Total Inflow (acre-feet)	Total Outflow (acre-feet)
STA-1E	2,400	4,600
STA-1W	8,900	12,000
STA-2	4,400	7,600
STA-3/4	16,000	13,000
STA-5/6	500	0

Water Depths
(4/12/2026)




Figure S-1. STA depths and flow volumes

0 CFS Lake release capacity in Eastern Flow Path:
 4/13/2026 to 4/19/2026
 Subject to change weekly as dry season progresses




Figure S-2. Eastern Flow Path Weekly Status Report

100 CFS Lake release capacity in Central Flow Path:
4/13/2026 to 4/19/2026
Subject to change weekly as dry season progresses




Figure S-3. Central Flow Path Weekly Status Report

0 CFS Lake release capacity in Western Flow Path:
4/13/2026 to 4/19/2026




Figure S-4. Western Flow Path Weekly Status Report

Basic Concepts and Definitions for STA Weekly Status Report

- **Inflow:** Sum of flow volume at all inflow structures to an STA.
- **Lake Inflow:** Portion of the STA total inflow volume that originates from Lake Okeechobee.
- **Outflow:** Sum of flow volume at outflow structures from an STA.
- **Total Phosphorus (TP):** Total mass of phosphorus in all its forms; including particulate, dissolved, etc.
- **Inflow Concentration:** TP concentration is the mass of TP in micrograms per liter of water, $\mu\text{g/L}$ or ppb. Inflow concentration refers to the flow-weighted mean TP from all inflow structures over a period of time.
- **Outflow Concentration:** The flow-weighted mean TP from all outflow structures over a period of time. The outflow concentration represents the reduction of inflow TP achieved by STA treatment of the inflow water.
- **WQBEL:** The STA outflow concentration that is required upon completion of the Restoration Strategies projects by December 2025. The outflow concentration shall not exceed 13 ppb as an annual flow weighted mean in more than 3 out of 5 water years on a rolling basis and shall not exceed 19 ppb as an annual flow weighted in any water year.
- **Flow-Way (FW):** One or more treatment cells connected in series. Cells typically have emergent aquatic vegetation (EAV) in the front portion of the flow-way followed by a mix of EAV and submerged aquatic vegetation (SAV)
- **Vegetation Status:** Healthy means the vegetation condition is good and will allow the STA to perform as designed. Stressed means the vegetation is showing signs of poor health, such as browning or areas of vegetation die-off, or the cell contains undesirable vegetation such as floating exotic vegetation requiring treatment. The TP reduction capability of the STA is affected when the vegetation condition is poor.
- **Phosphorus Loading Rate (PLR):** Mass of inflow TP in grams, divided by total treatment area of STA in square meters, per year. In general, a 365-day value of less than 1.0 is needed for an STA to perform optimally. A PLR of 2.0 is considered very high and a PLR of 3.0 is considered extremely high. The TP reduction capability of the STA is affected when the PLR is high, very high and extremely high.
- **Online:** Online status means the FW can receive and treat inflow.
- **Online with Restriction:** The FW can receive and treat inflow, but the amount of flow or water level may be limited temporarily. For example, a vegetation rehabilitation effort may require reduced flows through an area while the new plants are establishing, or nesting by protected species may require a certain water level not to be exceeded.
- **Offline:** The FW is unable to receive and treat inflow due to repairs, construction, or other prohibitive reasons.
- **Depth:** Difference between the average surface water level in a cell and the average ground elevation in that cell. Target depths, or depths between flow events, are between 1.25 ft to 1.5 ft. As depth approaches or drops below zero, an increasing percentage of the cell is considered dry and STA conditions deteriorate. An increase in depth above target depth is expected with increasing flow. However, as depth increases much above the target depth and is sustained over a period of time, it can be detrimental to vegetation health and overall STA treatment performance.
- **Note:** The data provided in this summary report were developed using a combination of provisional and quality-assured flow and water quality data. In some cases, best professional judgment was used to estimate missing data and revise questionable data. Values provided are not considered final but are appropriate for use in STA operational decision-making.

Everglades

Water Conservation Area

Regulation Schedules

WCA-1: Stage at the 1-8C gauge rose above the regulation line last week to 0.26 feet above the A1 Zone regulation line on Sunday, April 12, 2026 (**Figure EV-1**).

WCA-2A: Water depth at the 2-17 gauge increased last week and is now stabilizing. Stage was around 1.07 feet above the regulation line on Sunday (**Figure EV-2**).

WCA-3A: The 3-gauge average remains well within Zone B, continuing on a steady downward trend last week. On Sunday, stages were 1.7 feet NGVD29 below the Zone A regulation line. Stage at Gauge 62 (NW corner) resumed a downward trend away from the regulation line and was below the Upper Schedule line by 1.84 feet on Sunday (**Figures EV-3 and EV-4**).

Water Depths

The SFWDAT model output for April 12, 2026, illustrates a reversal in recessions within both WCA-1 and WCA-2A. WCA-1 increased in water depth with only a small area in the north now approaching ground level. WCA-2A also exhibited an increase in water depth over the past week with only a small portion of the north-central region now approaching ground level and still remaining slightly more ponded in the southeast. WCA-3A remains very dry, having received the least amount of rainfall in the system over the last month. In Everglades National Park (ENP), Taylor Slough, and potentially Shark River Slough, more hydrologic connection from north to south is helping flows go into Florida Bay. Comparing current conditions to water depths over the last twenty years, WCA-1 depths are now primarily above the 60th percentile and in the 80th percentile in the NE/SE portions of that basin. Southern WCA-2A depths are also at the 60th percentile in the central region and 80th percentile in the NE/SE. Big Cypress National Preserve depths are now above average in the center of the preserve. But even above average, depths for this time of year are below ground along Tamiami Trail. Within WCA-3A and WCA-3B, depths remain below the 10th percentile. Conditions in Southeastern and central ENP, remain above average. The southwest coastline of ENP is falling back down to below average depths. See **Figures EV-5 through EV-6**.

Taylor Slough and Florida Bay

Most stages decreased across Taylor Slough over the past week, with an average decrease of 0.11 feet for the week. Changes ranged from -0.50 feet at E112 in the northern slough to +0.13 feet at CT50R in the C-111 area (**Figure EV-7 and Figure EV-8**). Taylor Slough water levels are above the recent average (WY1993-2016) for this time of year by 8.0 inches compared to before the Florida Bay Initiative (starting in 2017), a decrease of 2.2 inches relative to last week. The stages at Craighead Pond (CP) and Taylor Slough Bridge (TSB) are below the estimated historical average by 0.11 and 0.80 feet, respectively.

Average salinity in Florida Bay was 31.5, a decrease of 1.1 from last week. Salinity changes ranged from -3.4 at Terrapin Bay (TB) in the central nearshore region to +1.3 at Johnson Key (JK) in the western region (**Figure EV-7**). Salinity remains above the estimated historical average and within the WY2001-2016 Interquartile Range (IQR) for all three regions. The Central region is now at the 25th percentile (**Figure EV-9**). Salinity in the western region remains near the hypersalinity threshold. Bay-wide salinity is above its recent average (WY1993-2016) for this time of year by 0.2, a decrease of 1.7 from last week.

Salinity at the Taylor River (TR) station in the mangrove zone (tracked for the Florida Bay MFL) was 1.5, a decrease of 0.9 from last week (**Figure EV-10**). The 365-day moving sum of flow from the five major creeks (McCormick Creek, Taylor River, Mud Creek, Trout Creek, West Highway Creek) was unable to be assessed due to missing data.

Average rainfall across Taylor Slough and Florida Bay was 0.75 inches over the past week, based on the 18 gauges used for this report. Rainfall ranged 0.02 inches at P37 in the southern slough to 2.72 inches at Highway Creek (HC) in the eastern nearshore region (**Figure EV-11**). Wind directions and speeds in Florida Bay ranged from 1.7 mph NW on April 7th to 26.4 mph NE on April 10th (**Figure EV-11**).

The Taylor River, Mud Creek, and West Highway Creek flow stations are currently offline until further notice, so data from all five major creeks are unable to be assessed. Based on the available data from Trout and McCormick Creeks, average daily flow totaled 988 ac-feet, with net positive flows for the week. Total daily creek flow ranged from -1,377 ac-feet on April 7th to 2,679 ac-feet on April 10th (**Figure EV-13**). Average daily flow from Alligator Creek was 13 ac-feet, with net positive flows for the week (**Figure EV-13**).

Implications/considerations for water management.

- Due to current drought conditions, slower recessions in WCA-3A can help protect the wetland ecology from damaging dry downs as the dry season continues.
 - Poor wading bird nesting is expected to continue for the fourth year into the 2026 nesting season.
 - With continued dry season drought conditions, conserving water within the WCAs, especially WCA-3A, will continue to be ecologically beneficial.
- Depths remain above average (~1.3 feet) in southern WCA-2A where shallower conditions on average are needed to recover ridge and slough habitat. However, holding water high in the north of the system, as further dry conditions are predicted, may prove ecologically beneficial as the Everglades dry down begins to accelerate due to increasing evapotranspiration. Input of water into areas prone to wildfire and peat oxidation may help to buffer the worst of the ecological stress.
- Freshwater input into Taylor Slough and the C-111 basin could help moderate salinities and support recovery of estuarine conditions in Florida Bay.

- Conserving water in the WCAs while providing freshwater input to the sloughs of ENP will require careful consideration of a balance between the upstream and downstream ecological needs of the system. Individual regional recommendations can be found in **Table EV-2**.

Table EV-2. Previous week’s rainfall and water depth changes in Everglades basins.

Everglades Region	Rainfall (inches)	Stage change (feet)
WCA-1	2.48	+0.19
WCA-2A	1.54	+0.13
WCA-2B	1.10	-0.06
WCA-3A	0.68	-0.08
WCA-3B	1.43	+0.16
ENP	0.84	-0.03




Figure EV-1. WCA-1 stage hydrographs and regulation schedule.




Figure EV-2. WCA-2A stage hydrographs and regulation schedule. $11.94 - 1.49 = 10.45, 1.05$




Figure EV-3. WCA-3A stage hydrographs (three-gauge average, 3-69W) and regulation schedule. $8.24 - 1.5 = 6.74, 1.68$




Figure EV-4. WCA-3A stage hydrograph (Deer gauge; Site 62) and regulation schedule.




Figure EV-5. Everglades water depths from two months ago (left), one month ago (center) and present (right), based on SFWDAT.




Figure EV-6. Present water depths (April 12, 2026) compared to the day of year relative to average (percentile) over the previous 20 years.




Figure EV-7. Taylor Slough water depths and Florida Bay salinities with changes since a week ago.




Figure EV-8. Taylor Slough water depth time series for Taylor Slough Bridge (TSB; northern slough) and Craighead Pond (CP; southern slough).




Figure EV-9. Eastern (top panel), Central (middle panel) and Western (bottom panel) Florida Bay daily average salinities with WY2001-2016 interquartile (25-75 percentile) ranges (IQR) and estimated historical daily average salinities. The hypersalinity threshold indicates the level at which salinities start to become harmful to seagrass.




Figure EV-10. Daily average salinity at Taylor River (TR) tracked for the Florida Bay MFL criteria. The 365-day total creek flow MFL metric is not currently available due to missing creek flow data.




Figure EV-11. Daily average rain across Taylor Slough and Florida Bay, along with hourly average wind speed and direction (measured at Long Key) in Florida Bay over the past four weeks.




Figure EV-12. Top: daily average creek flow summed between the five major creeks with estimated historical daily flow over the past four weeks (**note:** data from Taylor River, Mud Creek and West Highway Creek are currently unavailable since November 21st, 2025). Bottom: Daily average Alligator Creek flow data. N/A indicates missing data.

Table EV-2. Weekly water depth changes and water management recommendations

SFWMD Everglades Ecological Recommendations, April 12, 2026 (red is new)			
	Weekly change	Recommendation	Reasons
WCA-1	Stage increased by 0.19 feet.	A recession of no faster than 0.05 feet per week.	Conserve water, maintain within basin and downstream habitat and wildlife. Maintain maintenance access for vegetation management.
WCA-2A	Stage increased by 0.13 feet	A recession of no faster than 0.05 feet per week.	Maintain within basin (north versus south) and downstream habitat and wildlife.
WCA-2B	Stage decreased by 0.06 feet	A recession of no faster than 0.12 feet per week.	Protect within basin and downstream habitat and wildlife.
WCA-3A NE	Stage decreased by 0.01 feet	A recession of no faster than 0.05 feet per week.	Conserve water, maintain within basin and downstream habitat and wildlife. Provide suitable depths for aquatic prey and protect against peat soil loss during the dry season.
WCA-3A NW	Stage decreased by 0.01 feet	A recession of no faster than 0.05 feet per week.	
Central WCA-3A S	Stage decreased by 0.19 feet	A recession of no faster than 0.05 feet per week.	Conserve water, maintain within basin and downstream habitat and wildlife. Provide suitable depths for aquatic prey and protect against peat soil loss during the dry season.
Southern WCA-3A S	Stage decreased by 0.12 feet		
WCA-3B	Stage increased by 0.16 feet	A recession of no faster than 0.12 feet per week.	Protect within basin and downstream habitat and wildlife.
ENP-SRS	Stage decreased by 0.03 feet.	Make discharges to ENP according to COP protocol, considering up/down stream ecological conditions.	Protect within basin and upstream habitat and wildlife.
Taylor Slough	Stage changes ranged from -0.50 feet to +0.13 feet	Move water southward as possible.	When available, provide freshwater to promote water movement.
FB- Salinity	Salinity changes ranged from -3.4 to +1.3	Move water southward as possible.	When available, provide freshwater to promote water movement.